

Q : 755/755

Accuracy: - 0.00%

97

598

599

600

601

602

All

Q : 609 During chest auscultation a nurse will associate wheezing sound with

- A. Lung inflammation, loss of lubricating pleural fluid
- B. Fluid in the airways, opening of the collapsed alveoli
- C. Interstitial pneumonia, bronchitis
- D. Build up secretions, tumor

[Check Answer](#)[NEXT Q](#)[PREV](#)

Q : 755/755

Accuracy: 0.00%

97

598

599

600

601

602

All

Q:611 Causes of vesicovaginal fistula include:

- A. Hypertension, atherosclerosis
- B. Atherosclerosis, diabetes mellitus
- C. Obesity, pelvic inflammatory disease
- D. Cervical cancer, obstetric injury

[Check Answer](#)[NEXT Q&A](#)[RESET](#)[PREV Q&A](#)

Q : 755/755

Accuracy: - 0.00%

- 97
- 598
- 599
- 600
- 601
- 602
- All

Q:613 A complication of chronic bronchitis

- A. Pulmonary hypertension.
- B. Left ventricular heart failure.
- C. Status asthmaticus.
- D. Left lung collapse.

Check Answer

PREV QUE

RESET

NEXT QUE

← Q : 755/755

Accuracy:- 0.00%

- 97
- 598
- 599
- 600
- 601
- 602
- All

Q :617 Intra Aortic balloon pump therapy is used for treatment of:

- A. Congestive heart failure
- B. Cardiogenic shock
- C. Pulmonary edema.
- D. Aortic insufficiency

Check Answer

PREV QUE

RESET

NEXT QUE



Q : 755/755

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q:618 The most important health message to share with a family whose members has depression on medication is

- A. To Inform clinician immediately of the side effect!
- B. That 3 - 6 weeks must elapse before the effects are seen
- C. To give medication at the right time and dose
- D. To avoid stopping medication without medical advice

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)

Accuracy:- 0.00%



Q:619 The manoeuvre that involves assisting the mother on the hands and knees to increase the pelvic diameters as used in dystocia is the

- A. McRoberts Manoeuvre
- B. Gaskin Maneuver
- C. Zavanelli Manoeuvre
- D. Lovset Manoeuvre

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)

Q : 755/755

Accuracy:- 0.00%

- 597
- 598
- 599
- 600
- 601
- 602
- All

Q : 621 Bone marrow biopsy is required for diagnosis of.

- A. Hodgkin's lymphoma
- B. Megaloblastic anemia
- C. Thalassemia
- D. Thrombocytosis

Check Answer

PREV QUE

RESET

NEXT QUE



Q : 755/755

Accuracy: 0.00%

97

598

599

600

601

602

All

Q:620 Concerning the endocrine system changes in pregnancy, the following statement is incorrect

- A. Thyroid hormone production increases
- B. Insulin production increases thus increasing glucose available to the foetus
- C. Slight enlargement of the thyroid glands
- D. Levels of Oestrogen and Progesterone decrease

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)

Accuracy:- 0.00%

- 97
- 598
- 599
- 600
- 601
- 602
- All

Q:629 The factor that does NOT contribute to nonunion of the fracture is:

- A. Interposition of tissue between the bone ends
- B. Inadequate immobilisation that disrupts callus formation
- C. Excessive space between bone fragments
- D. Reduced skeletal muscle contractions of the extremities

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Accuracy:- 100%

97

598

599

600

601

602

Q:724 The following patients require reverse barrier Nursing

- A. Severe burns, premature baby
- B. Light for dates baby, cholera case
- C. Severe malnutrition, patient with ebola
- D. Severe burns, hepatitis

Check Answer

Accuracy:- 0.00%

- 97
- 598
- 599
- 600
- 601
- 602
- All

Q :628 Social learning that is based on pleasurable and exciting experiences is referred to as

- A. Positive socialisation
- B. Deliberate socialisation
- C. Planned socialisation
- D. Natural socialisation

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Q : 755/755



Accuracy:- 0.00%

97

598

599

600

601

602

All

Q : 705 Neuroleptic are:

- A. Major tranquillisers which block dopamine receptors thus produce calm effects.
- B. Minor tranquillisers with mild cam effects on neurotic patients.
- C. Mood stabilisers thus elevate the mood of a depressed patient.
- D. Selective serotonin reuptake inhibitors which block serotonin.

[Check Answer](#)



Q : 754/755

Accuracy:- 100%

97

598

599

600

601

Q :709 Depo-provera is a family planning method

- A. Contain progesterone and estrogen
- B. Helps in regulation of menstrual cycle
- C. Is contraindicated in breastfeeding women
- D. Contains progesterone only

Check Answer

Q : 755/755

Accuracy: 0.00%

- 597
- 599
- 600
- 601
- 602
- All

Q:703 The WHO standard case definition of a suspected case of Covid 19 Include:

- A. Acute onset of any three or more of the following signs or symptoms: Fever, cough, general weakness/fatigue, headache, myalgia, sore throat, coryza, dyspnoea, anorexia/nausea/vomiting, diarrhoea, altered mental status
- B. A patient who meet a clinical criteria above and is a contact of a probable or confirmed case, or linked to a COVID 19 cluster 3
- C. A person with recent onset of anosmia (loss of smell) or ageusia (loss of taste) in the absence of any other identified cause
- D. A person with a positive Nucleic Acid Amplification test (NAAT)

Check Answer

Q : 755/755

Accuracy 0.00%

597

598

599

600

601

602

All

Q : 702 Scientific management was advocated by:

- A. Max Weber.
- B. Fredrick Taylor
- C. Henry Fayol
- D. B F Skinner

[Check Answer](#)

Q : 755/755

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q:704 A 10 months old child with cough for the last two days,chest indrawing and 46 breaths per minute is classified to be having.

- A. Pneumonia.
- B. Very severe pneumonia.
- C. Febrile disease.
- D. Very severe Febrile disease.

[Check Answer](#)

Accuracy 0.00%

597

598

599

600

601

602

All

Q : 698 In management of severe dehydration, the total amount of fluid administered to a 6 month old child Include:

- A. 30mls/kg of ringer lactate in the 1st 30 minutes and 70ml/kg in the next 2 hours.
- B. 30mls/kg of ringer's lactate in the 1st 1 hour and 70mls/kg in the next 5 hours.
- C. 30mls/kg of half strength Darrows in the 1st 30 minutes and 70mls/kg in the next 2 hours.
- D. 30mls/kg of half strength Darrows in the first 1 hour and 70mls/kg in the next 5hours

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q:683 Water washed diseases are spread because there is.

- A. Lack of water protection.
- B. Water contamination.
- C. Lack of treatment.
- D. Inadequate water availability.

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Accuracy: 0.00%

597

598

599

600

601

602

All

Q : 689 The clinical assessment demonstrated in severe dehydration is:

- A. Skin turgor of less than 2 seconds.
- B. Sunken fontanelle
- C. Marked oliguria
- D. Decreased blood pressure

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)[Submit](#)

597

598

599

600

601

602

All

Q : 697 The nurse needs to carefully assess the complaint of pain of the elderly because old people.

- A. Are expected to experience chronic pain.
- B. Have a decreased pain threshold.
- C. Experience reduced sensory perception.
- D. Have altered mental function.

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Accuracy: 0.00%

97

598

599

600

601

602

All

Q : 691 Basal metabolic rate is the energy expended.

- A. When resting by not working and in fasting state.
- B. When doing moderate activities.
- C. To make food available to the body.
- D. When sleeping.

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)[Submit](#)

97

598

599

600

601

602

All

Q:693 The most common cause of pneumonia in children less than 5 years.

- A. *Streptococcus pneumoniae*.
- B. Group beta-hemolytic streptococcus.
- C. *Haemophilus influenzae*.
- D. *Staphylococcus aureus*.

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)[Submit](#)

Accuracy - 0.00%

97

598

599

600

601

602

All

Q : 699 The purpose of vaccine forecasting is:

- A. Ordering vaccine to avoid stock out
- B. Determining the population that require vaccination
- C. Estimating the number of persons immunised
- D. Maintenance of potency of vaccine

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)[Submit](#)

Q : 755/755

Accuracy: 0.00%

97

598

599

600

601

602

All

Q:692 In speech development, a child develops single monosyllabic words at the age of.

- A. 7-8 months
- B. 9 to 10 months first of
- C. 11 to 12 months.
- D. 12 to 14 months

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)[Submit](#)

Accuracy:- 0.00%

All

602

601

600

599

598

597

Q :681 Combined oral contraceptive

- A. Suppresses Ovulation
- B. Prohibit implantation of the egg.
- C. Makes cervical mucus hostile to sperm
- D. Acts as a barrier by destroying the sperms

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Q : 755/755

Accuracy: 0.00%

97

598

599

600

601

602

All

Q :674 Indication for using zinc sulphate in a child with diarrhoea is.

- A. Control diarrhoea.
- B. Reduces the severity of diarrhoea.
- C. Reduces duration of diarrhoea.
- D. Reduces frequency of diarrhoea.

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q : 680 Blood from a person infected with onchocerciasis cannot be used for diagnosis because.

- A. The onchocerca larvae take months to mature to adults.
- B. Onchocerca volvulus cannot be seen with a microscope.
- C. Microfilariae are found in fibrous nodules under the skin.
- D. Infective larvae are found under the skin.

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)[Submit](#)

Accuracy: 0.00%

All

597

598

599

600

601

602

Q : 684 Students are passive in the following teaching method.

- A. Practical.
- B. Lecture.
- C. Simulation.
- D. Project.

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q :682 The international nurses week is celebrated in the month of.

- A. March.
- B. April.
- C. May
- D. June

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Q : 755/755

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q :676 Insufficient nutritional quantities of vitamin D and phosphorus during childhood result in .

- A. Marasmus.
- B. Kwashiorkor.
- C. Osteomalacia
- D. Rickets.

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)

Q : 755/755

Accuracy - 0.00%

All

602

601

600

599

598

597

Q :679 The dietary sources of vitamin B12 include.

- A. Legumes and citrus fruits.
- B. Liver and green vegetables
- C. Fish and eggs.
- D. Milk and green

Check Answer

PREV QUE

RESET

NEXT QUE

Q : 755/755

Accuracy:- 0.00%

597

598

599

600

601

602

All

Q:671 The use of dapson in prophylaxis therapy for some HIV positive clients protects against.

- A. *Cryptococcus meningitis*.
- B. *Pneumocystis carinii pneumonia*.
- C. *Mycobacterium tuberculosis*.
- D. Bacterial pneumonia.

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)

Q : 755/755

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q : 672 During school health program

- A. All school children need in be served under one roof
- B. Groups need to be formed and served as per their need
- C. Teachers should be served first for them to able to control students
- D. Parent's teachers and students should be served together

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)

Accuracy:- 0.00%

- 598
- 599
- 600
- 601
- 602
- All

... legal document used for involuntary admission.

- A. MOH 614.
- B. MOH 637.
- C. MOH 613.
- D. MOH 638.

Check Answer

PREV QUE

RESET

NEXT QUE

Q : 755/755

Accuracy: 0.00%

598

599

600

601

602

All

Q : 669 The danger sign during pregnancy include:

- A. Severe headache
- B. Abdominal pain
- C. Nausea and vomiting.
- D. Vaginal discharge.

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)

Accuracy:- 0.00%

- 97
- 598
- 599
- 600
- 601
- 602
- All

Q :666 Psychomotor learning domain deals with.

- A. Attitude.
- B. Knowledge.
- C. Teaching methods
- D. Skills.

Check Answer

PREV QUE

RESET

NEXT QUE

Q : 755/755

Accuracy: - 0.00%

97

598

599

600

601

602

All

Q : 667 The major signs used in making diagnosis of rheumatic fever using Jones criteria is:

- A. Chorea, migratory polyarthritiis, arthralgia
- B. Chorea, carditiis . Erythema marginatum
- C. Chorea carditiis and fever
- D. Positive antistreptolysin , chorea, carditiis

[Check Answer](#)

Accuracy: 0.00%

All

597

598

599

600

601

602

Q : 665 The main purpose of identifying the syndrome in a client with an STI is to confirm

- A. Client's risk levels
- B. Presence of Sff
- C. Specific causative organism
- D. Flow chart instructions

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Accuracy - 0.00%

All

602

601

600

599

598

597

Q :665 The main purpose of identifying the syndrome in a client with an STI is to confirm

- A. Client's risk levels
- B. Presence of STI
- C. Specific causative organisms
- D. Flow chart instructions

Check Answer

PREV QUE

RESET

NEXT QUE

Q : 755/755

Accuracy: 0.00%

97

598

599

600

601

602

All

ncing of Maslow's hierarchy of human needs is.

- A. Physiological, belonging, safety, self-esteem, self-actualization.
- B. Self-actualization, safety, self-esteem, belonging, physiological.
- C. Physiological, safety, belonging, self-esteem, self-actualization.
- D. Safety, belonging, physiological, self-actualization, self-esteem.

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)

Q : 755/755

Accuracy:- 0.00%

- 97
- 598
- 599
- 600
- 601
- 602
- All

Q:664 The mortality rate that uses the estimated total mid-year population as the denominator is:

and always be

- A. Age-specific mortality rate
- B. Sex-specific mortality rate
- C. Crude mortality rate
- D. Cause-specific mortality rate

Check Answer

Q : 755/755

Accuracy:- 0.00%

- 97
- 598
- 599
- 600
- 601
- 602
- All

Q : 660 Maternal mortality ratio is the number of maternal deaths per

- A. 100,000 live births
- B. 100,000 women of reproductive age
- C. 10,000 women of reproductive age
- D. 10,000 live births

Check Answer

PREV QUE

RESET

NEXT QUE

Accuracy:- 0.00%

- 97
- 598
- 599
- 600
- 601
- 602
- All

Q:662 The MOST common cause of secondary amenorrhoea is almost always

- A. Drugs
- B. Nervous disorder
- C. Pregnancy
- D. Hormonal disturbance

Check Answer

PREV QUE

RESET

NEXT QUE

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q : 659 Breastfeeding promotes uterine involution because it,

- A. Stimulates the production of human placental lactogen that contracts the uterus
- B. Promotes secretion of prolactin To increase uterine contractions
- C. Causes the pituitary To secrete oxytocin to contract the uterus
- D. Emulates the secretion of prostaglandins to stimulate uterine contraction

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)

597

598

599

600

601

602

All

Q : 656 The route of drug administration with the highest bioavailability is the

- A. Sublingual
- B. Intramuscular
- C. Intravenous
- D. Rectal

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Q : 755/755

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q :658 The following is correct regarding sterilisation:

- A. Destruction of all microorganisms including bacterial spores
- B. Heat is the most effective agent for killing microorganisms
- C. Sterilization can use dry or moist heat
- D. Use of antiseptics in destroying microorganisms

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)

602

601

600

599

598

597

Q : 654 Paradoxical breathing is associated with the following condition.

- A. Lung contusions
- B. Chest wall lacerations
- C. Flail chest
- D. Pulmonary embolism

Check Answer

NEXT QUE

RESET

PREV QUE

Submit

Accuracy:- 0.00%

- 97
- 598
- 599
- 600
- 601
- 602
- All

Q:638 Delivery of the after coming head of the breech is accomplished through

- A. Burns-marshall maneuver
- B. Mauriceau-smellie veit maneuver
- C. Loveset maneuver
- D. McRoberts maneuver

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q :652 The best method of curtailing the second stage of labour in heart disease is by.

- A. Prophylactic forceps
- B. Prophylactic ventouse
- C. Spontaneous delivery with episiotomy
- D. Caesarean section

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)[Submit](#)

Accuracy:- 0.00%

- 597
- 598
- 599
- 600
- 601
- 602
- All

Q :636 Intense elation with feelings of Grandeur is:

- A. Euthymic mood
- B. Anhedonia
- C. Euphoria
- D. Alexithymia

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Q : 755/755

Accuracy:- 0.00%

- 97
- 598
- 599
- 600
- 601
- 602
- All

Q :632 In pustular psoriasis -

- A. An associated arthritis occurs in children
- B. Lesions are confined to the trunk
- C. Lesions are filled with pus
- D. There is always an eruption in the mouth.

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Q : 755/755

Accuracy: 0.00%

All

602

601

600

599

598

597

Q : 637 The diagonal conjugate is measurement between the following structures.

- A. Sacral promontory and coccyx
- B. Sacral promontory and superior edge of the pubic symphysis.
- C. Sacral promontory and inferior edge of the pubic symphysis.
- D. Coccyx and ischial spine.

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Q : 755/755



Accuracy: 0.00%

97

598

599

600

601

602

All

Q : 634 In a 29 year old female client who is being successfully treated for Cushing syndrome, a nurse would expect a decline in:

- A. Bone mineralization.
- B. Hair loss
- C. Menstrual flow
- D. Serum glucose

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)[Submit](#)

Q : 755/755

Accuracy: - 0.00%

- 597
- 598
- 599
- 600
- 601
- 602
- All

Q : 631 The medication associated with hearing loss as an adverse effect is:

- A. Gentamicin
- B. Fluconazole
- C. Amoxicillin
- D. Ceftriaxone

Check Answer

PREV QUE

RESET

NEXT QUE

Submit



Q : 755/755

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q :600 The main ion involved during muscle contraction is:

A. Calcium

B. Sodium.

C. Chloride

D. Potassium.

Check Answer

Q : 755/755

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q :633 The presenting diameters in a deflexed head are.

- A. Biparietal, sub occipitobregmatic.
- B. Bitemporal, sub occipitobregmatic.
- C. Biparietal, occipitofrontal.
- D. Occipitofrontal, suboccipital bregmatic.

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)[Submit](#)

Q : 755/755

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q :602 A mother has delivered the placenta. You note that the Shiny surface of the placenta was delivered first (i) The delivery mechanism that was used is; AND (ii) is this the maternal or fetal surface of the placenta?

- A. Matthew Duncan mechanism, maternal
- B. Schultze mechanism, maternal
- C. Schultze mechanism, fetus
- D. Matthew Duncan mechanism, foetus

[Check Answer](#)

Q : 755/755

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q:603 Nursing codeethics Is:

- A. The sum total of rules and regulations by which a profession is governed
- B. A written list of profession's values and standard of conduct
- C. The written and non-written moral principles in a given profession
- D. Norms and principles which should be observed by the professionals.

[Check Answer](#)

Q : 755/755

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q :601 Cardinal clinical features of Cholelithiasis include:

- A. High fever, anorexia, abdominal tenderness.
- B. Bleeding disorder, anorexia, weight loss
- C. Nausea, crampy pain, dark colored urine
- D. Greyish stool, jaundice, colic abdominal pain

[Check Answer](#)

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q:605 A 28 year old Para 3 Gravida 4 is admitted at 34 weeks with a history of mild per vaginal bleeding and abdominal pain. Her blood pressure is 140/90mmHg The likely diagnosis is:

- A. Placenta praevia
- B. Placenta abruption
- C. Preterm labour
- D. Urinary tract infection

[Check Answer](#)



Q : 755/755

Accuracy: - 0.00%

97

598

599

600

601

602

All

Q : 606 Laser ablation is indicated in treatment of:

- A. Cancer of the breast stage 0
- B. Stage III cancer of the ovary.
- C. Stage II cancer of the uterus.
- D. Stage 0 cancer of cervix.

[Check Answer](#)

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q : 604 Autoimmune diseases of Nervous system Includes:

- A. Encephalitis, Brain abscess
- B. Guillain barre syndrome, Meningitis
- C. Multiple sclerosis, Myasthenia Gravis
- D. Parkinson Disease, Alzheimer's Disease

[Check Answer](#)



Q : 755/755

Accuracy: - 0.00%

97

598

599

600

601

602

All

Q : 608 Oral hypoglycemic drugs are classified to:

- A. Metformin, glyburide, Glucovance
- B. Glipizide, Amaryl, glyburide.
- C. Micronase, Glucotrol, Prandin
- D. Nateglinide, Starlix, Glucophage

[Check Answer](#)[NEXT QUE](#)[PROCT](#)

Q : 755755

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q:607 The priority nursing intervention for a mother with prolapsed pulsating cord include

- A. Cover the exposed cord with a sterile dressing
- B. Administer oxygen to the mother
- C. Place the mother in knee chest or exaggerated trendelenburg position
- D. Return the cord into the vaginal canal

[Check Answer](#)[RESET](#)[NEXT Q](#)



Q : 755/755

Accuracy: 0.00%

97

598

599

600

601

602

All

Q :612 Uterine sub - involution is indicated by:

- A. Bulky uterus.
- B. Constant uterine size for several days
- C. Tender uterus
- D. Mild fever

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)



Q : 755/755

Accuracy: 0.00%

97

598

599

600

601

602

All

Q : 614 Penetrating chest trauma commonly results in which pathologic mechanisms -

- A. Hypoxemia, ciliary function is reduced, fibrosis
- B. Hypovolemia, Hypoxemia, Cardiac failure
- C. Fibrosis, alveolar dilatation, Hypovolemia
- D. Cardiac failure, ~~mycobacterial~~ infection, atelectasis

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)

Q : 755/755

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q:615 The features a midwife would find on examination of the bregma in a newborn include

- A. Diamond shaped, pulsations, Large membranous space
- B. Large membranous space, lies along coronal suture, pulsations
- C. Diamond shaped, lies at the junction of frontal suture and coronal suture, pulsations
- D. Does not pulsate, triangular shaped, larger membranous space

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)



Q : 755/755

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q : 610 Microorganisms responsible for hospital acquired pneumonia are-

- A. Streptococcus pneumonia, haemophilus influenza, legionella pneumophila
- B. Chlamydia trachomatis, mycobacterium tuberculosis, parainfluenza virus
- C. Pseudomonas aeruginosa, staphylococcus aureus, klebsiella pneumonia
- D. Pneumocystis carinii, cytomegalovirus, aspergillus fumigatus

[Check Answer](#)[RESET](#)[NEXT Q](#)

Q : 755/755

Accuracy: 0.00%

- 597
- 598
- 599
- 600
- 601
- 602
- All

Q :622 The following is suggestive of low hemoglobin

- A. Jaundice.
- B. Cyanosis
- C. Plethora
- D. Pallor

Check Answer

PREV QUE

RESET

NEXT QUE



Q : 755/755

Accuracy: - 0.00%

97

598

599

600

601

602

All

Q : 623 Direct cause of maternal Death.

- A. Anaemia, unsafe abortions
- B. HIV/AIDS, sepsis
- C. Pre eclampsia, antepartum hemorrhage.
- D. Malaria, eclampsia

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)



Q : 755/755

Accuracy - 0.00%

597

598

599

600

601

602

All

Q:616 An early sign of obstructed labour is

- A. Slow descent
- B. Dehydration
- C. Bandl's ring
- D. Hot and dry vagina

Check Answer

PREV QUE

RESET

NEXT QUE



Q : 755/755

Accuracy: 0.00%

97

598

599

600

601

602

All

Q : 625 Clinical manifestation of whooping cough during the catarrhal stage is

- A. Cough
- B. Vomiting
- C. Whoop
- D. Sneezing

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)[Submit](#)

Accuracy:- 0.00%

- 597
- 598
- 599
- 600
- 601
- 602
- All

Q:626 In relation to waste management, human bodytissues are classified as:

- A. Clinical waste
- B. Pathological waste
- C. Pharmaceutical waste
- D. Radiation waste

Check Answer

PREV QUE

RESET

NEXT QUE

Submit



Q : 755/755

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q :624 Effective hazard management largely rely on:

- A. Government agencies
- B. Emergency responses.
- C. Pre disaster planning.
- D. Volcanoes

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Accuracy:- 100%

97

598

599

600

601

602

AI

Q:725 The immediate intervention in sequence a midwife will undertake to manage primary postpartum haemorrhage includes

- A. Call for help, stop bleeding, resuscitate the mother
- B. Stop bleeding, resuscitate the mother, empty the uterus
- C. Rub up a contraction. give oxytocin, empty the uterus
- D. Administer oxytocin, call for medical aid, resuscitate the mother

[Check Answer](#)



Accuracy:- 100%

97

598

599

600

601

602

Q:723 The human sperms mature in the

- A. Vas deferens
- B. Spermatic cord
- C. Urethra
- D. Epididymis

[Check Answer](#)

Accuracy:- 100%

97

598

599

600

601

602

A

Q:719 A ten-year-old girl develops convulsions while in the ward. The priority nursing intervention to be prevented by the nurse is.

- A. To ensure safety of the patient.
- B. To ensure the Airway of the patient is open.
- C. To stop convulsions by administering diazepam.
- D. To put the spoon into the mouth of the patient to prevent biting of the third.

[Check Answer](#)

Accuracy:- 100%

97

598

599

600

601

602

Q:722 Nurse J is providing nursing care to a 5 year old boy with asthma. The desired clinical outcome for respiratory distress from asthma is

- A. The child remains asleep for 8 hrs at night.
- B. Child is in no pain or discomfort
- C. Airway remain patent
- D. Oral mucous membranes remain moist and pink

[Check Answer](#)

Accuracy:- 100%

97

598

599

600

601

602

All

Q:716 DNA polymerase chain reaction test for HIV exposed infants is usually done at.

- A. 6 weeks, if negative, repeat at 6-months and 12-months.
- B. 6 weeks, if negative, repeat at 3-months and 6-months.
- C. 6 weeks, if negative, repeat at 6-months and 9-months.
- D. 6 weeks, if negative, repeat at 9-months and 12-months.

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)

Q : 754/755

Accuracy:- 100%

97

598

599

600

601

602

All

Q:714 The activity that is only done during the first focused antenatal care visit is:

- A. Check on individual birth plans.
- B. Comprehensive history taking.
- C. Give iron and folate.
- D. Listen to foetal heart sounds.

[Check Answer](#)

Accuracy:- 100%

97

598

599

600

601

602

Q:710 A child with fever and any general danger sign should be classified as having.

- A. Uncomplicated malaria.
- B. Acute ear infection.
- C. Very severe Febrile disease.
- D. Mastoiditis.

[Check Answer](#)

597

598

599

600

601

602

All

Q:701 The recommended management of meningitis in a child include

- A. Benzyl penicillin 50,000IU/kg for 14 days
- B. Benzyl Penicillin 100,000IU/kg for 14 days
- C. Ceftriaxone 50 mg/kg for 7 days
- D. Ceftriaxone 100 mg/kg for 14 days

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Accuracy:- 0.00%

- 97
- 598
- 599
- 600
- 601
- 602
- All

Q:661 The main cause of primary hypothyroidism is -

- A. Autoimmune destruction of the thyroid cells
- B. Resistance to the effects of thyroid hormones
- C. Loss of thyroid hormones due to kidney disease
- D. Hypothalamus disorders.

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)

Accuracy:- 100%

97

598

599

600

601

602

All

Q:715 For the client who is using oral contraceptives, the nurse will inform the client to take the pill at the same time each Day to accomplish.

- A. Decrease the incidence of nausea.
- B. Maintain hormone levels.
- C. Reduce side effects.
- D. Prevent drug interactions

[Check Answer](#)

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q:653 A 32 yr old primigravida comes to the labor ward for induction. Which drug is given for cervical ripening

- A. Salbutamol
- B. Prostaglandin E2
- C. Syntocinon
- D. Hyoscine bromide

[Check Answer](#)[PREV QUE](#)[NEXT QUE](#)[RESET](#)[Submit](#)



Q : 755/755



Accuracy: 0.00%

97

598

599

600

601

602

All

Q:650 A 35 year old Para 4 Gravida 5 presents to the ANC and a diagnosis of anaemia is made. The likely cause of anemia in her case is

- A. Folate deficiency
- B. Sickle cell anaemia
- C. Iron deficiency anaemia
- D. pernicious anemia

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)[Submit](#)

97

598

599

600

601

602

All

Q :655 General care during postpartum period includes assessment of

- A. Breasts, abdomen, fundus
- B. Breasts, abdomen, muscular skeletal system
- C. Breasts, abdomen, neurological responses
- D. Breasts, abdomen, congenital abnormality

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Accuracy: 0.00%

All

597

598

599

600

601

602

Q : 651 The signs of hypoglycemia in a full-term baby include.

- A. High-pitched cry, irritability, roving eyes.
- B. Poor feeding, convulsions, jitteriness.
- C. Rigidity of the trunk, apneic episodes, convulsions
- D. Twitching, high-pitched cry, apneic episodes

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Accuracy:- 0.00%

597

598

600

601

602

All

Q : 657 The modifiable risk factors to coronary artery disease are;

- A. Advanced age, obesity, smoking
- B. Family history, advanced age, smoking
- C. Hypertension, diabetes, smoking
- D. Obesity, family history, smoking

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

- 597
- 598
- 599
- 600
- 601
- 602
- All

Q : 646 Characteristic of first degree burns include?

- A. Erythema and painful
- B. Chilling and dry
- C. Leathery and painless
- D. Black and painful.

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Q : 755/755

Accuracy:- 0.00%

- 97
- 598
- 599
- 600
- 601
- 602
- All

Q :647 MR. T, a 67 year old man has smoked for the last 20 years and worked as a Quarrymen for 30 years. the following is a less significant risk for lung cancer

- A. Age
- B. Lifestyle
- C. Gender
- D. Occupation.

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Q : 755/755

Accuracy:- 0.00%

All

602

601

600

599

598

97

Q :648 The purpose of emergency assessment is to

- A. Make a diagnostic conclusion
- B. Delegate nursing responsibility
- C. Teach the client about his or her health
- D. Establish a database concerning the client

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Q : 755/755

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q : 649 When performing routine assessment on a mother post delivery, the midwife notes that the uterus is soft and displaced to the left of the umbilicus. The immediate plan of action will be to.

- A. Perform fundal massage and assist the patient to the bathroom.
- B. Notify the obstetrician immediately.
- C. Continue to monitor the mother as this is normal finding post delivery.
- D. Administer 10 international units of syntocinon.

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)[Submit](#)

Q : 755/755

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q :644 Characteristics of cephalohematoma include

- A. Pits on pressure, tends to enlarge, crosses a suture line
- B. Pits on pressure, circumscribed swelling, does not cross a suture line
- C. Does not pit on pressure, crosses a suture line, circumscribed swelling
- D. Does not pit on pressure, does not cross a suture line, circumscribed swelling

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)[Submit](#)

Accuracy: 0.00%

97

598

599

600

601

602

All

Q : 640 Early signs intracranial pressure include,

- A. Decreased heart rate, decrease in level of consciousness, restlessness
- B. Decreased heart rate, systolic blood pressure increase, agitation
- C. Increased heart rate, decrease in level of consciousness, lethargy
- D. Increased heart rate, systolic blood pressure increase, stupor

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)[Submit](#)



Accuracy:- 0.00%

97

598

599

600

601

602

All

Q :643 The most effective method of physically assessing for left ventricular function in a patient with left ventricular failure is:

- A. Listening to heart sounds.
- B. Assessment of peripheral and sacral edema.
- C. Assessment of jugular venous distention.
- D. Monitoring for organomegaly.

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)[Submit](#)

Q : 755/755

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q:645 In Choanal atresia.

- A. There is incomplete canalization of the GIT thus altered nutritional state
- B. A web of tissues occludes the nasal passage thus altered breathing pattern
- C. It is possible to insert a nasogastric tube but occlusion usually occurs immediately
- D. There is a harsh sound produced during breathing due to constricted passages

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)[Submit](#)

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q : 639 Common causes of secondary amenorrhea includes:-

- A. Pelvic tumours, excess prostaglandins, chronic PID
- B. Cervical stenosis, excess prostaglandins, oral contraceptive pills.
- C. Ischemia of the uterus, narrow cervical canal, endometriosis
- D. Uterine fibroid, chronic PID, endometriosis

[Check Answer](#)[PREV QUE](#)[RESET](#)[NEXT QUE](#)[Submit](#)

Accuracy:- 0.00%

97

598

599

600

601

602

All

Q :642 The dosage for vitamin A supplementation for children at 6 months is

- A. 50,000 IU
- B. 100,000 IU
- C. 150,000 IU
- D. 200,000 IU

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

Accuracy:- 0.00%

- 97
- 598
- 599
- 600
- 601
- 602
- All

Q :635 Haematological disorder caused by deficiency of factor VIII is

- A. Sickle cell disease.
- B. Hemophilia.
- C. Leukemia.
- D. Polycythemia.

Check Answer

PREV QUE

RESET

NEXT QUE

Submit

597

598

599

600

601

602

All

Q :641 The common cause of oligohydramnios >•

- A. Esophageal atresia
- B. Placenta hemangioma
- C. Renal agenesis
- D. Diabetes mellitus

Check Answer

PREV QUE

RESET

NEXT QUE

Submit