

REPRODUCTIVE HEALTH E-MCQS

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
1	22-04-2022	Winny	CM	Voi	19	34.5%	F	abdominal circumference	all of the above	weak pulse	dyspnea	manual vacuum aspiration	plunger	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262843cfb7f576b1aee8b54
2	22-04-2022	Wayua	Nursing	Kmtc port reitz	31	56.4%	F	abdominal circumference	all of the above	weak pulse	dyspnea	misoprostol	cannula	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/62628497629f256b61d51546
3	22-04-2022	Jaycee	Nursing	Voi	6	10.9%	M	femur length	plan for childbirth	weak pulse	wheezing	misoprostol	syringe	the wall of the uterus feel smooth	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/626284b47cf5316af6388049
4	22-04-2022	Yooh	Clinical med	Meru	30	54.5%	M	crown rump length	all of the above	hypertension	dyspnea	awaiting spontaneous expansion only on oxytocin	nostle	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262850f6aa11b6aa934d210

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
5	22-04-2022	Velinta	Nursing	Mosoriot	25	45.5%	F	abdominal circumference	all of the above	rapid pulse	dyspnea	manual vacuum aspiration	nostle	the wall of the uterus feel smooth	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262855ea86a606b5bba2525
6	22-04-2022	Jou	Co	Nakuru	18	32.7%	M	crown rump length	all of the above	weak pulse	dyspnea	manual vacuum aspiration	syringe	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/6262855f7cf5316af638812e
7	22-04-2022	Joan cherotich	Nursing	Kmtc mosoriot	25	45.5%	F	abdominal circumference	all of the above	rapid pulse	dyspnea	manual vacuum aspiration	nostle	the wall of the uterus feel smooth	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626285671a974c6b370a3679
8	22-04-2022	F.P	Bsc Clinical Medicine	Uzima university	18	32.7%	M	crown rump length	all of the above	anaemia	dyspnea	awaiting spontaneous expansion only on oxytocin	cannula	the vacuum in the syringe decreases	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262857c1a974c6b370a36e0
9	22-04-2022	Martin Njeru	Clinical med	Meru uni	42	76.4%	M	biparietal diameter	all of the above	hypertension	dyspnea	awaiting spontaneous expansion only on oxytocin	nostle	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/6262858925aeec6af0449a8c

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
10	22-04-2022	Imani	Clinical medicine	Gluk	25	45.5%	F	chest circumference	all of the above	raised bilirubin	dyspnea	misoprostol	syringe	when the patient feels min pain	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/626285bb02e67c6b8884e245
11	22-04-2022	Collins Rollins	Nursing	Mosoriot	24	43.6%	M	abdominal circumference	all of the above	weak pulse	spider naevi	awaiting spontaneous expansion only on oxytocin	nostle	the wall of the uterus feel smooth	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626285d56aa11b6aa934d25f
12	22-04-2022	Eliud	Cm	Km	36	65.5%	M	femur length	all of the above	raised bilirubin	dyspnea	awaiting spontaneous expansion only on oxytocin	syringe	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/626285ffa04146a8767dfa7
13	22-04-2022	Haska	Med	Jkuat	18	32.7%	M	femur length	identify health education and counseling needs	raised bilirubin	dyspnea	misoprostol	nostle	when the patient feels min pain	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262862b8a13dd6b3d495f5c

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
14	22-04-2022	Ramli	Nursing	Nakuru	0	0%	M	chest circumference	identify existing problems	rapid pulse	rhonchi	Ergometrine	plunger	the vacuum in the syringe decreases	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/6262866aa056e36ab0a300b1
15	22-04-2022	John	Mbchb	Mku	30	54.5%	M	abdominal circumference	all of the above	raised bilirubin	dyspnea	misoprostol	syringe	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/62628672da70dc6acca72a1c
16	22-04-2022	Bensoul	Cm	Bsa	24	43.6%	M	chest circumference	all of the above	weak pulse	dyspnea	misoprostol	syringe	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/626286ec02e67c6b8884e2bf
17	22-04-2022	Frances	Nursing	Kmtc	24	43.6%	M	biparietal diameter	all of the above	weak pulse	dyspnea	manual vacuum aspiration	cannula	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262872c8a13dd6b3d495faa
18	22-04-2022	Mohaa	Nursing	W	25	45.5%	F	abdominal circumference	identify existing problems	hypertension	dyspnea	manual vacuum aspiration	syringe	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/626287378a13dd6b3d495fad

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
19	22-04-2022	Mogire	Nursing	Kabarnet	18	32.7%	M	crown rump length	plan for childbirth	raised bilirubin	dyspnea	Ergometrine	nostle	the uterus relaxes	a clot form within 7 minutes	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626287541a974c6b370a371d
20	22-04-2022	Ronoh Kipkoech	Orthopaedic	Makindu	36	65.5%	M	abdominal circumference	all of the above	raised bilirubin	dyspnea	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262877402e67c6b8884e307
21	22-04-2022	Justice	Clinical med	Ecuadro	31	56.4%	F	biparietal diameter	all of the above	raised bilirubin	microscopic hematuria	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626287917cf5316af6388163
22	22-04-2022	Esther	Orthopaedic trauma medicine	Mbooni	19	34.5%	F	femur length	all of the above	rapid pulse	dyspnea	Ergometrine	valve	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/6262881a02e67c6b8884e365
23	22-04-2022	Kevin mwangi	Orthopaedic	Goshen Medical	24	43.6%	M	abdominal circumference	all of the above	raised bilirubin	microscopic hematuria	misoprostol	valve	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262881dfa04146a8767e00f

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
24	22-04-2022	Edwin kolil	Community health nursing	Mosoriot campus	42	76.4%	M	crown rump length	all of the above	raised bilirubin	dyspnea	awaiting spontaneous expansion only on oxytocin	nostle	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262886f7cf5316af6388186
25	22-04-2022	Joseph	Clinical medicine	Kendu Adventist school of medical science	25	45.5%	F	crown rump length	all of the above	hypertension	dyspnea	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62628889a86a606b5bba25ae
26	22-04-2022	Diana Adhiambo	BSC clinical medicine	EGERTON	37	67.3%	F	femur length	all of the above	raised bilirubin	dyspnea	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/626288b68a13dd6b3d495fd4
27	22-04-2022	Sam	Cmed	Must	24	43.6%	M	femur length	all of the above	rapid pulse	dyspnea	misoprostol	nostle	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626289776aa11b6aa934d2fb

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
28	22-04-2022	Vivian Kosgei	Clinical medicine	Iten kmtc	31	56.4%	F	femur length	all of the above	rapid pulse	dyspnea	awaiting spontaneous expansion only on oxytocin	nostle	the uterus relaxes	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/62628a04fb7f576b1aee8cd2
29	22-04-2022	Yvonne Atieno	Clinical medicine	Bomet	19	34.5%	F	femur length	all of the above	rapid pulse	microscopic hematuria	manual vacuum aspiration	cannula	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/62628a0c8723736a8d5081e7
30	22-04-2022	Cornelius Matende	Clinical Medicine	Kmtc	30	54.5%	M	abdominal circumference	all of the above	raised bilirubin	microscopic hematuria	awaiting spontaneous expansion only on oxytocin	nostle	red or pink form but no more tissue is aspirated	a clot form within 7 minutes	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62628a6bda70dc6acca72ae4
31	22-04-2022	Edith Nakoba	Clinical medicine	Gluk	19	34.5%	F	femur length	all of the above	anaemia	dyspnea	misoprostol	syringe	red or pink form but no more tissue is aspirated	a clot form within 7 minutes	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/62628a74fb7f576b1aee8d5a
32	22-04-2022	Caroline	Clinical medicine	Nairobi	25	45.5%	F	abdominal circumference	all of the above	rapid pulse	dyspnea	manual vacuum aspiration	nostle	the vacuum in the syringe decreases	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62628a7a6aa11b6aa934d349
33	22-04-2022	Hosea Mokuwa Anyong'a	Diploma in community health nursing	Kapkatet campus	12	21.8%	M	abdominal circumference	all of the above	anaemia	rhonchi	misoprostol	nostle	the uterus relaxes	a clot form within 7 minutes	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/62628b077cf5316af638825f

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
34	22-04-2022	Mm	Bsc clin med	Egerton	31	56.4%	F	abdominal circumference	all of the above	weak pulse	microscopic hematuria	awaiting spontaneous expansion only on oxytocin	nostle	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/62628b388a13dd6b3d496100
35	22-04-2022	Robinson	Bsc Clinical medicine	Great lakes	36	65.5%	M	biparietal diameter	all of the above	raised bilirubin	dyspnea	awaiting spontaneous expansion only on oxytocin	cannula	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62628b8c1a974c6b370a38f9
36	22-04-2022	James	Krchn	kmtc	12	21.8%	M	crown rump length	all of the above	weak pulse	microscopic hematuria	Ergometrine	cannula	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62628ba9a056e36ab0a30222
37	22-04-2022	Laban mbau	Bsc clinical medicine	University of kabianga	36	65.5%	M	biparietal diameter	all of the above	hypertension	dyspnea	misoprostol	nostle	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/62628c0b8723736a8d508223

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
38	22-04-2022	Bee	CM	Me	25	45.5%	F	abdominal circumference	identify existing problems	raised bilirubin	dyspnea	manual vacuum aspiration	syringe	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62628c49a86a606b5bba2604
39	22-04-2022	Maryline	Nursing	Bomet	7	12.7%	F	crown rump length	identify health education and counseling needs	hypertension	dyspnea	manual vacuum aspiration	syringe	the uterus relaxes	a clot form within 7 minutes	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62628cba7eb0016ad32cc25e
40	22-04-2022	Sharon Chemutai	Bsn clinical medicine	Uzima University	43	78.2%	F	femur length	all of the above	raised bilirubin	dyspnea	awaiting spontaneous expansion only on oxytocin	nostle	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/62628d07629f256b61d5170c
41	22-04-2022	Sammy	CHN	Kisumu	12	21.8%	M	crown rump length	all of the above	anaemia	spider naevi	misoprostol	cannula	the vacuum in the syringe decreases	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/62628d4aa86a606b5bba2622
42	22-04-2022	BNO	MD	KP	24	43.6%	M	abdominal circumference	identify existing problems	raised bilirubin	microscopic hematuria	misoprostol	syringe	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/62628d94fb7f576b1aee8e8a

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
43	22-04-2022	Maryann	Nursing	Mkd	31	56.4%	F	crown rump length	all of the above	raised bilirubin	dyspnea	misoprostol	cannula	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/62628e87da70dc6acca72bae
44	22-04-2022	Alicia	Nursing diploma	Kitui	13	23.6%	F	abdominal circumference	all of the above	anaemia	wheezing	awaiting spontaneous expansion only on oxytocin	No Answer	No Answer	No Answer	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62628f78712cd06b82006f9e
45	22-04-2022	Dante Dante	Garissa	Garissa	13	23.6%	F	abdominal circumference	all of the above	rapid pulse	dyspnea	misoprostol	syringe	the uterus relaxes	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62628ff48a13dd6b3d496301
46	22-04-2022	Victor	Nursing	Kitale	6	10.9%	M	crown rump length	all of the above	weak pulse	microscopic hematuria	Ergometrine	syringe	when the patient feels min pain	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626290117cf5316af638837b
47	22-04-2022	Bee Mel	Nursing	Karatina	18	32.7%	M	crown rump length	all of the above	rapid pulse	dyspnea	misoprostol	cannula	the uterus relaxes	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262902f8a13dd6b3d49630a

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
48	22-04-2022	Duncan Ian	Mwingi	Mwingi	42	76.4%	M	biparietal diameter	all of the above	raised bilirubin	dyspnea	awaiting spontaneous expansion only on oxytocin	nostle	red or pink form but no more tissue is aspirated	a clot form within 7 minutes	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626290a7da70dc6acca72cc5
49	22-04-2022	Alex Thuku Kaminja	Clinical medicine and surgery	Mosoriot	31	56.4%	F	biparietal diameter	all of the above	hypertension	spider naevi	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626290b18723736a8d508314
50	22-04-2022	Ian charo	Cm	Msa	6	10.9%	M	abdominal circumference	all of the above	rapid pulse	spider naevi	manual vacuum aspiration	cannula	the uterus relaxes	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62629109629f256b61d51ad8
51	22-04-2022	Nicholas	Cm	Makindu	24	43.6%	M	abdominal circumference	all of the above	weak pulse	spider naevi	awaiting spontaneous expansion only on oxytocin	valve	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262911ada70dc6acca72cd3
52	22-04-2022	Mercy margaret	Diploma in Clinical medicine	Mku	25	45.5%	F	abdominal circumference	all of the above	rapid pulse	dyspnea	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/626291437eb0016ad32cc3ae

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
53	22-04-2022	Kinyanga	Cm	Kitui	12	21.8%	M	abdominal circumference	all of the above	hypertension	spider naevi	manual vacuum aspiration	cannula	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62629143da70dc6acca72cd7
54	22-04-2022	Eric	Bsc Clinical medicine	Uok	36	65.5%	M	biparietal diameter	all of the above	weak pulse	dyspnea	misoprostol	nostle	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262914a712cd06b82006fcf
55	22-04-2022	Dickie	D/CM	Kape	24	43.6%	M	abdominal circumference	all of the above	hypertension	dyspnea	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62629235da70dc6acca72ced
56	22-04-2022	Cynthia	Nursing	Lodwar	7	12.7%	F	abdominal circumference	all of the above	anaemia	microscopic hematuria	misoprostol	cannula	the wall of the uterus feel smooth	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/62629253629f256b61d51afb

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
57	22-04-2022	Titus langat	Medicine	Tenwek	30	54.5%	M	femur length	all of the above	raised bilirubin	spider naevi	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626292aefb7f576b1aee9031
58	22-04-2022	Tingeet	Clinical medicine	Nairobi	18	32.7%	M	crown rump length	identify health education and counseling needs	raised bilirubin	dyspnea	misoprostol	cannula	the uterus relaxes	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626292d5629f256b61d51b31
59	22-04-2022	mary	medicine	uon	13	23.6%	F	abdominal circumference	all of the above	rapid pulse	microscopic hematuria	manual vacuum aspiration	syringe	red or pink form but no more tissue is aspirated	a clot form within 7 minutes	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626292ef8723736a8d50851a
60	22-04-2022	Dorcas Njoroge	Clinical medicine	Egerton	25	45.5%	F	femur length	all of the above	hypertension	dyspnea	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62629417a056e36ab0a304f1
61	22-04-2022	Eliud isaiah	Cm	Kisii	6	10.9%	M	abdominal circumference	all of the above	weak pulse	spider naevi	misoprostol	syringe	the uterus relaxes	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626294818a13dd6b3d496543

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
62	22-04-2022	Samuel	KRN	Mathare	12	21.8%	M	chest circumference	all of the above	weak pulse	dyspnea	misoprostol	cannula	the uterus relaxes	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626296c9fb7f576b1aee90f0
63	22-04-2022	Rk	Med	Un	42	76.4%	M	femur length	all of the above	hypertension	dyspnea	awaiting spontaneous expansion only on oxytocin	nostle	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/62629781a056e36ab0a305dd
64	22-04-2022	Njiriri	Bachelor in Clinical Medicine	Egerton University	37	67.3%	F	femur length	all of the above	raised bilirubin	dyspnea	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/62629945da70dc6acca73015
65	22-04-2022	Lyntondan	Clinical medicine	Nyamira	13	23.6%	F	crown rump length	all of the above	hypertension	microscopic hematuria	manual vacuum aspiration	syringe	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62629967b8c93c6b14378816
66	22-04-2022	Karma Romeo	BSC clinical medicine	Meru university	30	54.5%	M	abdominal circumference	all of the above	rapid pulse	dyspnea	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62629b848a13dd6b3d496663

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
67	22-04-2022	Jully	Nursing	Nairobi	7	12.7%	F	abdominal circumference	identify existing problems	weak pulse	microscopic hematuria	manual vacuum aspiration	nostle	the vacuum in the syringe decreases	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62629c44fb7f576b1aee9227
68	22-04-2022	Susan	Nursing	Muranga	13	23.6%	F	abdominal circumference	identify health education and counseling needs	anaemia	dyspnea	misoprostol	syringe	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62629de325aee6af044a5f2
69	22-04-2022	James	Nursing	Siaya	24	43.6%	M	abdominal circumference	all of the above	hypertension	dyspnea	misoprostol	cannula	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/62629eba7cf5316af6388a2a
70	22-04-2022	Isaac Benson	Clinical medicine	Great Lakes	24	43.6%	M	crown rump length	all of the above	rapid pulse	dyspnea	misoprostol	syringe	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/62629f381a974c6b370a4699
71	22-04-2022	Sharline	Community health nursing	Kitale campus	13	23.6%	F	biparietal diameter	plan for childbirth	rapid pulse	rhonchi	Ergometrine	cannula	the vacuum in the syringe decreases	a clot form within 7 minutes	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/62629f691a974c6b370a469c

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
72	22-04-2022	Barno	Bsn	Nai	18	32.7%	M	crown rump length	all of the above	weak pulse	microscopic hematuria	misoprostol	nostle	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262a11fa056e36ab0a3084f
73	22-04-2022	Diana Odongo	Nursing	Mosoriot	31	56.4%	F	femur length	all of the above	raised bilirubin	microscopic hematuria	Ergometrine	syringe	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262a20d7eb0016ad32ccac4
74	22-04-2022	LAS	DCM	Nyamira	24	43.6%	M	abdominal circumference	all of the above	raised bilirubin	dyspnea	manual vacuum aspiration	cannula	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/6262a2bafa04146a8767ea3e
75	22-04-2022	Betty chepkurui	Clinical medicine	Aic litein MTC	19	34.5%	F	abdominal circumference	identify existing problems	raised bilirubin	spider naevi	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262a32d7eb0016ad32ccad2
76	22-04-2022	Vvvvv	Ccc	Gg	43	78.2%	F	crown rump length	all of the above	raised bilirubin	dyspnea	awaiting spontaneous expansion only on oxytocin	nostle	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262a3cc25aeec6af044a789

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
77	22-04-2022	Rosemary onyanggo	Nursing	Kaplong	13	23.6%	F	crown rump length	identify existing problems	raised bilirubin	spider naevi	manual vacuum aspiration	cannula	the uterus relaxes	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262a5998a13dd6b3d496823
78	22-04-2022	Ann	C.Medicine	Kemu	25	45.5%	F	crown rump length	all of the above	rapid pulse	wheezing	misoprostol	nostle	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262a6e88a13dd6b3d496841
79	22-04-2022	Husna	Clinical medicine and surgery	Thika	19	34.5%	F	chest circumference	identify health education and counseling needs	raised bilirubin	dyspnea	misoprostol	syringe	red or pink form but no more tissue is aspirated	a clot form within 7 minutes	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262a6f4a86a606b5bba2ca1
80	22-04-2022	Manuu	Nursing	Nyahururu	18	32.7%	M	abdominal circumference	all of the above	rapid pulse	dyspnea	Ergometrine	cannula	the uterus relaxes	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262a746fb7f576b1aee94da
81	22-04-2022	John Emmanuel	Clinical medicine	GREAT LAKES UNIVERSITY OF KISUMU	36	65.5%	M	biparietal diameter	all of the above	raised bilirubin	dyspnea	manual vacuum aspiration	syringe	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262a75f7cf5316af6388bc0

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
82	22-04-2022	Meshack	Bachelors of science in clinical medicine	Great Lakes University of Kisumu	24	43.6%	M	crown rump length	all of the above	rapid pulse	dyspnea	Ergometrine	nostle	the uterus relaxes	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262a97aa86a606b5bba2e0b
83	22-04-2022	Toni Ngikito	Cm	Kakamega	18	32.7%	M	abdominal circumference	all of the above	weak pulse	dyspnea	manual vacuum aspiration	cannula	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262aac9b8c93c6b14378dae
84	22-04-2022	Salim	Cm	Wby	30	54.5%	M	crown rump length	all of the above	raised bilirubin	dyspnea	misoprostol	nostle	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262aad725aee6af044a849
85	22-04-2022	Sam	C med	Must	37	67.3%	F	femur length	all of the above	raised bilirubin	dyspnea	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262ac92712cd06b82007591
86	22-04-2022	Nickdee	Bcm	Kisumu	42	76.4%	M	biparietal diameter	all of the above	hypertension	dyspnea	misoprostol	nostle	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262acf2fb7f576b1aee96f0

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
87	22-04-2022	Kimatu Mwanja	Clinical medicine and surgery	Machakos	30	54.5%	M	biparietal diameter	all of the above	anaemia	dyspnea	misoprostol	syringe	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262b153b8c93c6b1437910a
88	22-04-2022	Vinic	ECHN	Bomet	18	32.7%	M	biparietal diameter	all of the above	weak pulse	spider naevi	misoprostol	cannula	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/6262b2858a13dd6b3d496c78
89	22-04-2022	Faith chepngeno	Clinical medicine	GLUK	19	34.5%	F	crown rump length	all of the above	weak pulse	dyspnea	Ergometrine	cannula	the uterus relaxes	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262b451a056e36ab0a30d57
90	22-04-2022	Evans	Dcm	Kapkatet	18	32.7%	M	abdominal circumference	identify health education and counseling needs	raised bilirubin	spider naevi	misoprostol	valve	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262b6c1fb7f576b1aee9c7c
91	22-04-2022	Rik	Clinical medicine	University of kabianga	12	21.8%	M	crown rump length	all of the above	anaemia	spider naevi	manual vacuum aspiration	cannula	red or pink form but no more tissue is aspirated	a clot form within 7 minutes	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262bc4aa86a606b5bba35c8

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
92	22-04-2022	Enock Ngeno	Krchn	Matibabu	18	32.7%	M	crown rump length	all of the above	anaemia	dyspnea	misoprostol	cannula	the wall of the uterus feel smooth	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262be9b8723736a8d50928b
93	22-04-2022	Sharon	Clinical medicine	Kmtc	25	45.5%	F	biparietal diameter	all of the above	raised bilirubin	wheezing	manual vacuum aspiration	syringe	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262c7856aa11b6aa934e732
94	22-04-2022	Chemu	Bsm	Egerton	31	56.4%	F	abdominal circumference	all of the above	raised bilirubin	dyspnea	misoprostol	nostle	red or pink form but no more tissue is aspirated	a clot form within 7 minutes	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262ccd58a13dd6b3d497321
95	22-04-2022	Winfer	Clinical medicine	Jkuat	31	56.4%	F	abdominal circumference	all of the above	raised bilirubin	dyspnea	manual vacuum aspiration	syringe	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262cd14a056e36ab0a315c1
96	22-04-2022	Chol	Clinical Medicine	Kabarak university	19	34.5%	F	abdominal circumference	all of the above	hypertension	spider naevi	misoprostol	nostle	the vacuum in the syringe decreases	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262ced07eb0016ad32cdc2e

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
97	22-04-2022	Cecilia kavee	Nursing	Makindu	31	56.4%	F	crown rump length	all of the above	raised bilirubin	dyspnea	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262cf19a86a606b5bba39aa
98	22-04-2022	Bb	Nursing	MAKINDU	7	12.7%	F	abdominal circumference	identify existing problems	rapid pulse	dyspnea	misoprostol	cannula	when the patient feels min pain	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262d1f7629f256b61d53573
99	22-04-2022	Hassan	Nursing	Mkd	25	45.5%	F	abdominal circumference	all of the above	rapid pulse	dyspnea	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	a clot form within 7 minutes	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/6262d2bea86a606b5bba3a77
100	22-04-2022	Lucy	Krchn	0701858338	30	54.5%	M	biparietal diameter	all of the above	rapid pulse	dyspnea	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	a clot form within 7 minutes	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262d358fa04146a8767fe1b
101	22-04-2022	Robert	nursing	bomet	30	54.5%	M	abdominal circumference	all of the above	raised bilirubin	microscopic hematuria	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262d37c02e67c6b8884f961

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
102	22-04-2022	Halima Dama	Clinical medicine	Msambweni	25	45.5%	F	biparietal diameter	all of the above	raised bilirubin	dyspnea	misoprostol	syringe	when the patient feels min pain	a clot form within 7 minutes	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/6262d4bb8723736a8d509aae
103	22-04-2022	David	DM	NAI	30	54.5%	M	crown rump length	all of the above	hypertension	dyspnea	manual vacuum aspiration	syringe	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262d748da70dc6acca745c6
104	22-04-2022	Abdi gza	Clinical medicine	Garissa	42	76.4%	M	femur length	all of the above	raised bilirubin	dyspnea	awaiting spontaneous expansion only on oxytocin	nostle	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262d9a76aa11b6aa934ecdf
105	22-04-2022	Ruto Rogony	Orthotrauma	Kaptumo	12	21.8%	M	abdominal circumference	identify existing problems	rapid pulse	dyspnea	misoprostol	valve	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262da9f02e67c6b8884fb53
106	22-04-2022	Abdiii	Nursing	Garisa	30	54.5%	M	abdominal circumference	all of the above	hypertension	dyspnea	Ergometrine	cannula	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262db248a13dd6b3d4976d5

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
107	22-04-2022	Celine Molly	Orthotrauma	Kaptumo	49	89.1%	F	biparietal diameter	all of the above	raised bilirubin	dyspnea	awaiting spontaneous expansion only on oxytocin	nostle	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262db7225aeec6af044bfc2
108	22-04-2022	Wycliffe lokichu	Nursing	Msambweni	24	43.6%	M	abdominal circumference	all of the above	hypertension	dyspnea	awaiting spontaneous expansion only on oxytocin	valve	the vacuum in the syringe decreases	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262db8225aeec6af044bfc4
109	22-04-2022	Nurdin kulow Ali	Clinical medicine	Garissa	48	87.3%	M	biparietal diameter	all of the above	raised bilirubin	dyspnea	awaiting spontaneous expansion only on oxytocin	nostle	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262dc6fa056e36ab0a31e38
110	22-04-2022	Elizabeth mutunga	Clinical medicine	Makindu	31	56.4%	F	crown rump length	all of the above	rapid pulse	dyspnea	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262dd5b1a974c6b370a5f41

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
111	22-04-2022	Asenath Kariuki	Cm	Matuga	19	34.5%	F	biparietal diameter	all of the above	rapid pulse	spider naevi	manual vacuum aspiration	cannula	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/6262dd5d629f256b61d53a4c
112	22-04-2022	Jacob Owaga Karewa	Diploma in registered nursing and midwifery	Homabay	24	43.6%	M	chest circumference	all of the above	rapid pulse	spider naevi	Ergometrine	nostle	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262dfc1fb7f576b1aeeb04e
113	22-04-2022	Ramaa	Nursing	Kitui	19	34.5%	F	biparietal diameter	identify existing problems	weak pulse	dyspnea	Ergometrine	cannula	when the patient feels min pain	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/6262e14d6aa11b6aa934f11d
114	22-04-2022	Ramla	Nursing	Thika	19	34.5%	F	biparietal diameter	all of the above	anaemia	microscopic hematuria	misoprostol	nostle	the vacuum in the syringe decreases	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/6262e2fb1a974c6b370a61a6

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
115	22-04-2022	Wilfred	Nursing	Kisii	6	10.9%	M	femur length	all of the above	anaemia	wheezing	manual vacuum aspiration	cannula	the uterus relaxes	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/6262e7a525aee6af044c236
116	22-04-2022	Gladys	Cm	Kmtc	37	67.3%	F	crown rump length	all of the above	raised bilirubin	dyspnea	misoprostol	nostle	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262e8237cf5316af638a4b2
117	22-04-2022	Miryam cherop	Clinical medicine	Kitale	25	45.5%	F	abdominal circumference	all of the above	rapid pulse	dyspnea	manual vacuum aspiration	cannula	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/6262e8d68723736a8d509cba
118	22-04-2022	Zephaniah kiprop	Orthopedic	Sigowet	12	21.8%	M	abdominal circumference	identify health education and counseling needs	raised bilirubin	microscopic hematuria	manual vacuum aspiration	nostle	when the patient feels min pain	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/6262eb88a056e36ab0a3247e
119	22-04-2022	Christine	CM	Kapee	13	23.6%	F	abdominal circumference	all of the above	weak pulse	microscopic hematuria	misoprostol	nostle	the vacuum in the syringe decreases	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262ec49fb7f576b1aeeb287

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
120	22-04-2022	Robert	Nursing	Kilifi	30	54.5%	M	abdominal circumference	all of the above	raised bilirubin	dyspnea	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262ecbc7cf5316af638a537
121	22-04-2022	Me	T	T	24	43.6%	M	femur length	all of the above	hypertension	dyspnea	manual vacuum aspiration	nostle	the uterus relaxes	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262ee4bfb7f576b1aeeb2a7
122	22-04-2022	Aden	Nursing	Embu	12	21.8%	M	chest circumference	all of the above	weak pulse	dyspnea	misoprostol	cannula	the uterus relaxes	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262eed3712cd06b8200949b
123	22-04-2022	Moses	clinical medicine	nairobi	30	54.5%	M	biparietal diameter	all of the above	hypertension	spider naevi	awaiting spontaneous expansion only on oxytocin	nostle	the uterus relaxes	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262f0cc712cd06b8200952b

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
124	22-04-2022	Alpha	Nursing	Gatundu	13	23.6%	F	crown rump length	all of the above	anaemia	dyspnea	Ergometrine	cannula	the wall of the uterus feel smooth	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262f24ffb7f576b1aeeb2e8
125	22-04-2022	Judy peter	Nursing	Makindu	19	34.5%	F	abdominal circumference	all of the above	anaemia	dyspnea	manual vacuum aspiration	valve	the wall of the uterus feel smooth	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262f62625aeec6af044c445
126	22-04-2022	Asha	Nursing	Bome	19	34.5%	F	biparietal diameter	all of the above	weak pulse	dyspnea	Ergometrine	cannula	the vacuum in the syringe decreases	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262fa191a974c6b370a65fa
127	22-04-2022	Diana	Cm	Km	31	56.4%	F	biparietal diameter	all of the above	weak pulse	dyspnea	misoprostol	nostle	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262fa2c8a13dd6b3d497b40
128	22-04-2022	Yvonne	Cm	Nakuru	31	56.4%	F	crown rump length	all of the above	weak pulse	dyspnea	awaiting spontaneous expansion only on oxytocin	cannula	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262faa7da70dc6acca7508c

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
129	22-04-2022	Beatrice	KRCHN	Nyabondo MTC	37	67.3%	F	abdominal circumference	all of the above	raised bilirubin	dyspnea	awaiting spontaneous expansion only on oxytocin	cannula	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6262ff2ca056e36ab0a3261a
130	22-04-2022	Unknown	Unknown	Unknown	19	34.5%	F	abdominal circumference	all of the above	anaemia	microscopic hematuria	manual vacuum aspiration	nostle	when the patient feels min pain	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6262ff898a13dd6b3d498029
131	22-04-2022	Jebet	Orthopaedic	Kakamega	31	56.4%	F	abdominal circumference	all of the above	raised bilirubin	dyspnea	misoprostol	cannula	the uterus relaxes	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/626301a602e67c6b888504b2
132	22-04-2022	Charles	Nursing	Kmtc	1	1.8%	F	abdominal circumference	identify health education and counseling needs	hypertension	microscopic hematuria	manual vacuum aspiration	cannula	the wall of the uterus feel smooth	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62630cc4fb7f576b1aeeb9d9
133	22-04-2022	Francis	Clinical medicine and minor surgery	Mkd	24	43.6%	M	biparietal diameter	identify existing problems	hypertension	microscopic hematuria	manual vacuum aspiration	nostle	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626311d56aa11b6aa934f6f6

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
134	22-04-2022	Roidah Kwamboka Arasa	Clinical medicine	Nakuru	25	45.5%	F	biparietal diameter	identify existing problems	raised bilirubin	dyspnea	manual vacuum aspiration	cannula	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62631457b8c93c6b1437ac31
135	23-04-2022	Hellen Achieng	clinical medicine	nyamira	19	34.5%	F	abdominal circumference	all of the above	weak pulse	dyspnea	manual vacuum aspiration	syringe	the uterus relaxes	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/6263837402e67c6b888515c7
136	23-04-2022	Ajevi liz	Clinical medicine	Makindu	31	56.4%	F	biparietal diameter	all of the above	raised bilirubin	microscopic hematuria	awaiting spontaneous expansion only on oxytocin	cannula	red or pink form but no more tissue is aspirated	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626386bafa04146a87681e3b
137	23-04-2022	Collins	Clinical medicine	Makindu kmtc	19	34.5%	F	crown rump length	all of the above	raised bilirubin	microscopic hematuria	misoprostol	cannula	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626387e61a974c6b370a7bd3

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
138	23-04-2022	Caxton	Nursing	Gatundu	24	43.6%	M	biparietal diameter	all of the above	rapid pulse	dyspnea	Ergometrine	nostle	the wall of the uterus feel smooth	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626389a4b8c93c6b1437c3e4
139	23-04-2022	Masayi florida	diploma in clinical medicine	St marys	19	34.5%	F	biparietal diameter	all of the above	rapid pulse	dyspnea	misoprostol	cannula	when the patient feels min pain	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62639e916aa11b6aa9350d6d
140	23-04-2022	Robai Linet	Bsc Clinical Medicine and community Health	Meru University	43	78.2%	F	femur length	all of the above	raised bilirubin	dyspnea	awaiting spontaneous expansion only on oxytocin	valve	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6263b291712cd06b8200bc2f
141	23-04-2022	Wycliff Nyaanga	Nursing	Nairobi	18	32.7%	M	abdominal circumference	identify existing problems	hypertension	dyspnea	misoprostol	cannula	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6263b4e1629f256b61d55f6d
142	23-04-2022	Emelda	Clinical medicine	Kapkatet campus	25	45.5%	F	femur length	all of the above	rapid pulse	dyspnea	manual vacuum aspiration	nostle	the vacuum in the syringe decreases	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/6263d1dafb7f576b1aeceb33

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
143	23-04-2022	Faith Nyatichi	Nursing	Nakuru kmtc	25	45.5%	F	abdominal circumference	all of the above	hypertension	dyspnea	manual vacuum aspiration	cannula	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6263d5c47eb0016ad32d1ff0
144	23-04-2022	Joyce nasambu	Bcm	Uzima university	25	45.5%	F	crown rump length	all of the above	hypertension	dyspnea	awaiting spontaneous expansion only on oxytocin	syringe	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6263d939fa04146a87684258
145	23-04-2022	Tracy	Orthopaedic trauma medicine	Sigowet	31	56.4%	F	abdominal circumference	all of the above	hypertension	dyspnea	Ergometrine	cannula	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/6263e59ca056e36ab0a36c21
146	23-04-2022	Job	CM	Kapkatet	0	0%	M	abdominal circumference	plan for childbirth	anaemia	wheezing	manual vacuum aspiration	syringe	the wall of the uterus feel smooth	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/6263f220fb7f576b1aeef582
147	23-04-2022	Phrunk	Nursing	MSU	24	43.6%	M	biparietal diameter	all of the above	anaemia	dyspnea	Ergometrine	cannula	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and a contracted uterus	https://quizzory.in/answer-sheet/626410128a13dd6b3d49ca24

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
148	23-04-2022	Phenny	Clinical medicine	Kisumu	13	23.6%	F	biparietal diameter	all of the above	rapid pulse	microscopic hematuria	manual vacuum aspiration	valve	when the patient feels min pain	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62641415da70dc6acca79e4e
149	23-04-2022	Sally	KRCHN	Makindu	25	45.5%	F	crown rump length	all of the above	rapid pulse	dyspnea	manual vacuum aspiration	cannula	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	a complete placenta and contracted uterus	https://quizzory.in/answer-sheet/626414e1a056e36ab0a38286
150	23-04-2022	Collen	Nursing	Maseno	6	10.9%	M	abdominal circumference	all of the above	weak pulse	microscopic hematuria	manual vacuum aspiration	plunger	the wall of the uterus feel smooth	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626415067cf5316af6390182
151	23-04-2022	Brian	Clinical Medicine	Siaya	12	21.8%	M	crown rump length	identify existing problems	rapid pulse	microscopic hematuria	misoprostol	syringe	red or pink form but no more tissue is aspirated	a clot failed to form after 7 minutes with a soft clot forms that breaks down easily	a complete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62641d2cb8c93c6b1437f843

S.NO	Submit Date	Name	course	campus	Total Marks (55)	Result	1. GENDER	2. the preferred fetal dimensions for estimating gestational age at 36 to 38 weeks is	3. the information obtained from the antenatal history can help the provider	4. a woman who suffers shock as a result of an obstetric emergency may have which of the following except	5. a pregnant woman has a severe anaemia typically present with	6. inevitable abortion at 18 weeks is best managed with	7. which of the following is not a component of manual vacuum aspiration kit	8. the MVA procedure is complete when	9. disseminated intravascular coagulopathy is suspected following when bedside clotting test is done if	10. tears of the cervix,vagina or perineum should be suspected when there is immediate postpartum hemorrhage and	Answer Sheet Link
152	23-04-2022	Lucas	Med	Maseno City campus	37	67.3%	F	abdominal circumference	all of the above	raised bilirubin	dyspnea	awaiting spontaneous expansion only on oxytocin	nostle	red or pink form but no more tissue is aspirated	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62641d2e8723736a8d510a00
153	23-04-2022	Nelima	Nursing	Nursing	19	34.5%	F	abdominal circumference	all of the above	raised bilirubin	dyspnea	misoprostol	syringe	the uterus relaxes	a clot fails to form within seven minutes when calcium gluconate is added to a 3 ml test tube of blo	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62641ff925aee6af045190b
154	23-04-2022	F	Nursing	Embu	19	34.5%	F	abdominal circumference	all of the above	weak pulse	dyspnea	misoprostol	nostle	the vacuum in the syringe decreases	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/626433ce629f256b61d58cbb
155	23-04-2022	George	Nursing	Maseno	12	21.8%	M	femur length	all of the above	rapid pulse	dyspnea	Ergometrine	cannula	when the patient feels min pain	bleeding fails to stop within seven minutes from my 1mm incision made on the inner aspect of a forea	an incomplete placenta and an atonic uterus	https://quizzory.in/answer-sheet/62643a9d1a974c6b370ab70b