

Dawn School of Medicine

UNIVERSITY OF NAIROBI
UNIVERSITY EXAMINATIONS 2015/2016

**EXAMINATIONS FOR THE DEGREE OF BACHELOR OF MEDICINE AND
BACHELOR OF SURGERY**

HPS 400: PSYCHIATRY

MULTIPLE CHOICE QUESTIONS

DATE: JUNE 28, 2016

TIME: 2.00 P.M. - 4.30 P.M.

INSTRUCTIONS:

1. This paper contains 100 multiple choice questions.
2. Only one Answer is correct.
3. Select the Best Answer out of the 5 choices.
4. Mark the correct answer on the answer sheet provided.
5. Follow the instructions on the answer sheet and do not tamper with it.

1. According to DSMIV the following are anxiety disorders except

- (a) Posttraumatic stress disorder - Trauma & Stressor related disorder
~~(b)~~ Conversion disorder - somatic
(c) Obsessive compulsive disorder - obsessive compulsive disorder
(d) Substance induced anxiety disorders ✓
(e) Anxiety disorder due to hyperthyroidism ✓

DSMIV

Generalized
Agoraphobia
Social anxiety
Selective mutism
Specific phobia

Panic anxiety
Substance / medication induced
Panic disorder

2. The key features in diagnosing posttraumatic stress disorder includes the following except

- A (a) History of significant traumatic event ✓ Stressor
~~(b)~~ Brain injury
~~(c)~~ Persistent Hyperarousal - difficulties sleeping, irritability, difficulty concentrating, overly alert, easily startled
B (d) Re-experiencing of the trauma - intrusive thoughts, flashbacks, intense emotional / physical upset at reminders,
C (e) Avoidance of reminders - Avoid thoughts / feelings, avoid activities, situations, places, can't recall important aspects
Loss of interest in activities, detached from others.

3. In Treatment of posttraumatic stress disorder the following are predictors of good prognosis except:

- (a) Rapid onset of symptoms
(b) Short duration of symptoms
(c) Strong social support
(d) Good premorbid functioning
~~(e)~~ Presence of schizophrenia in the patient

- ~~before ECT~~
- for depn
- ~~xmitter depn~~
- tonic inhib.
- in at responding a
- val
- i patient psychos
4. Clinical features of generalized anxiety disorders include the following except:
- Emotional numbing - PTN
 - Anxiety
 - Automatic hyperactivity - restlessness of breath, excessive sweating, palpitations, tachycardia, etc. cumm: ANS, PNS
 - Cognitive vigilance - irritability, easily startled, difficulty concentrating, mind going blank, sleep disturbance
 - Motor tension - shakiness, restlessness, headaches

5. Regarding electroconvulsive therapy, the following are true except:

- (a) May be used in treatment of manic episodes

- (b) It is safe treatment for depressed elderly patients unless there are contraindications

- (c) It is safe treatment for all pregnant mothers

- (d) It has been in use since 1930's 1938

- (e) It requires short acting anaesthesia (methohexitane or ms relaxant)

- hypnotic
ketamine

6. It is risky to conduct electroconvulsive therapy on patients with conditions listed below except:

- (a) Controlled hypertension & PT uncontrolled HT

- (b) Cerebral aneurysm

- (c) Space occupying lesion

- (d) Cerebral oedema

- (e) Recent myocardial infarction

7. In electroconvulsive therapy the following is true

- (a) It was discovered by BF Skinner

- (b) Temporary memory loss may occur

- (c) It is one of the dangerous psychiatric treatments

- (d) It is a pharmacotherapy

- (e) Bone fracture is one of the side effects

8. Signs of alcohol dependence includes the following except:

- (a) Tolerance

- (b) Withdrawal syndrome

- (c) Improved priority setting

- (d) Stereotyped pattern of use

- (e) Relieve drinking

9. The manifestation of Wernicke-Korsakoff syndrome is basically due to:

- (a) Brain damage

- (b) Thiamine deficiency

- (c) Dementia

- (d) Vitamin C deficiency

- (e) Deranged renal functioning

Symptoms (Clinical)

Ocular disturbance (ophthalmoplegia)

Changes in mental state (confusion)

Unsteady gait & gait (ataxia)

Others: stupor, hypotension, tachycardia,

hypothermia, epileptic seizures,

progressive hearing loss



The following are depot antipsychotic preparations except:

- (a) Fluphenazine decanoate ✓
(b) Haloperidol decanoate ✓
(c) Olanzapine pamoate ✓ psychotic antiepileptic. Causes metabolic synd.
(d) Zuclopentixol decanoate ✓
(e) Clozapine decanoate

Paliperidone palmitate
Risperidone microspheres ✓

Paliperidone palmate
Flupenthixol decanoate
Zyprexa risperidone

64.

Regression of milestones is a common presentation of childhood disorders. Which of the following medical conditions is associated with regression?

- (a) Central Nervous System infections
(b) Phenylketonuria
(c) Hypothyroidism
(d) Seizure disorder
(e) All of the above

65.

HIV/AIDS is a risk factor for all of the following disorders except:

- (a) Schizophrenia ✓
(b) Alcohol use disorder ✓
(c) Depression ✓ - major depressive disorder
(d) Adjustment disorder ✓
(e) Anxiety disorder ✓ - GAD, panic attacks

o somatization
o disturbance of conduct
o Delirium
o Suicide
o substance use

o Bipolar affective disorder.
o HIV assoc. Neurocognitive disorders (HANB)
o HIV assoc. dementia (HAD)
Minor Cognitive Motor system (MCMS)
→ assoc. with met. encephalopathy
C viral load > 3000 8.5x more likely)

66.

In addition to clinical and family history, a near confirmatory diagnostic test for a psychiatric disorder is:

- (a) Magnetic Resonance Imaging
(b) Single photon emission computed tomography
(c) Electroencephalogram
(d) Blood dopamine level
(e) None

67.

In the Four P's Model of psychiatric formulation, which domain of psychopathology is concerned with features that make the presenting condition endure and determine the severity of the condition or compliance issues?

- (a) Predisposing
(b) Precipitating
(c) Perpetuating
(d) Protective
(e) Persecutory

68.

What is the strongest risk factor for youth suicide?

- (a) Major depression ✓
(b) Bipolar disorder ✓
(c) Substance use disorder ✓
(d) Prior history of a suicide attempt ✓
(e) Borderline personality disorder

(59)

With regards to the parents of the child being evaluated, a child psychiatrist should take all of the following considerations except:

- (a) The expectations as well as feelings about the evaluation should be elicited from the child's parents
 - (b) Parents may bring a mixed heritage of guilt, shame, concern and fear of being judged in seeking help for their child
 - (c) Parents may have unrealistically optimistic fantasies of "absolution" of unconscious guilt or hopes of quick cures
 - (d) Parents generally inform the child of the nature of the evaluation prior to arrival
- ~~(e)~~ Parents generally have a well-formed understanding of psychiatric assessment and treatment from mass media sources

(60)

In which of the following situations would the risk of developing Schizophrenia be the lowest for a child?

- (a) A child's Schizophrenic parent 12% has a non Schizophrenic identical twin 0%
 - (b) A child's non Schizophrenic parent 0% has a Schizophrenic identical twin 4%
 - (c) A child's Schizophrenic parent 12% has a non Schizophrenic fraternal twin 0%
 - (d) A child's non Schizophrenic parent 0% has a Schizophrenic fraternal twin 4%
- ~~(e)~~ A child's non-Schizophrenic parent 0% has a non Schizophrenic sibling 0%

(61)

The following are correct about children below the age of 7 years in relation to understanding death in the family except:

- (a) Cannot comprehend the irreversibility of death
- (b) Can express their feelings in play therapy
- (c) Has limited ability to verbalize their feelings
- (d) The children tend towards magical thinking and ego centricity
- ~~(e)~~ The children are sensitive about being 'different' from their peers

(62)

In attention deficit hyperactivity disorder (ADHD), which of the following is NOT classified as a problem relating to poor attention?

- (a) Appears routinely forgetful
- ~~(b)~~ Experiences difficulties in waiting in turn
- (c) Dislikes tasks involving mental effort e.g. homework
- (d) Easily distracted
- (e) Appears not to listen, when spoken to

74.

Which ONE of the following was the first effective somatic therapy for Schizophrenia or bipolar disorder?

- (a) Chlorpromazine
- (b) ECT
- (c) Hypnotherapy
- (d) Insulin coma therapy
- (e) Pyrotherapy

75.

Which ONE of the following antidepressants is most suitable in a patient who is suffering from depression and has recently also had myocardial infarction?

- (a) Amitriptylline TCA
 - (b) Duloxetine SNRI
 - (c) Mirtazapine
 - (d) Paroxetine SSRI
 - (e) Sertraline SSRI
- CAB
Centraline
Citalopram
also fluoxetine

76.

Which of the following neurotransmitters promotes sleep?

- (a) Acetylcholine
- (b) Dopamine
- (c) Histamine
- (d) Noradrenaline
- (e) Serotonin

77.

Which ONE of the following can be used to reduce alcohol consumption in patients who are still consuming alcohol?

- (a) Acamprosate → craving in abstinent pts.
- (b) Bupropion - tx of nicotine addiction (cravings & withdrawal sympt)
- (c) Disulfiram - interferes with alcohol metabolism, ↑ acetaldehyde = headache, flushing, nausea
- (d) Nalmefene blocking serotonin, lack cardio & anxiety.
- (e) Naltrexone - blocks effects of opiates (precipitates withdrawal sympt)

78.

About the use of antipsychotics in the management of bipolar disorder, which of the following induces a metabolic syndrome?

- (a) Lamotrigine
- (b) Lithium ↑ gain
- (c) Lorazepam ↑ hyperglycaemia
- (d) Olanzapine ↑ hyperlipidemia
- (e) Sodium valproate ↑ wt

79.

In bipolar disorder, which of the following is useful in acute mania?

- (a) Bupropion
- (b) Carbamazepine & valproate
- (c) Imipramine
- (d) Sertraline
- (e) Tranylcypromine

Lamotrigine - bipolar depn esp Hx of

Carbamazepine - used in non-response.

body centered therapy let looks at & connects mind & body & uses both psychotherapy & healing therapy A healthy life, good exercise, marriage, dance

(69)

What is a simple alternative to the bell and pad method for the treatment of enuresis?

- (a) Waking up the child using a bull horn
- (b) Setting an alarm clock after 2 to 3 hours of sleep when the bladder may be reaching full capacity
- (c) Keeping the lights on at night to aid arousal when the bladder signals full capacity
- (d) Restricting fluids after 8pm
- (e) None of the above

(70)

The psychiatrist assessment of children commonly differs from the assessment of adults in ALL of the following ways except:

- (a) The child's behaviour may cause greater distress to the adults than to the child
- (b) Children may not recognize their behaviour as problematic to others
- (c) Children often seek out psychiatric assessment or treatment for themselves
- (d) Children may attribute problems to others and be unwilling to accept their own personal contribution to the problem
- (e) The adult's expectation for the child may exceed the child's abilities

(71)

Which of the following symptoms of ADHD (Attention Deficit Hyperactivity Disorder) is most likely to be outgrown when a child matures to adulthood?

- (a) Inattention
- (b) Hyperactivity
- (c) Impulsivity
- (d) Distractibility
- (e) Both hyperactivity and impulsivity

72.

A 74 year old man presents with memory impairment and behaviour that is out of character. Which of the following cognitive tests is appropriate to test frontal lobe function?

- (a) Benton Visual retention test
- (b) National adult reading test
- (c) Rey-Osterrieth Complex figure test
- (d) Rorschach ink blot test
- (e) Wisconsin Card Sorting test

73.

A 76 year old woman suffers a stroke due to a ruptured berry aneurysm in the anterior cerebral artery and the parietal lobe is affected. Which ONE of the following tests would you expect to be abnormal?

- (a) Colour vision testing
- (b) Copying intersecting pentagons
- (c) Luria's motor test
- (d) Semantic memory
- (e) Three object recall

80.

Which of the following statements is true concerning lithium?

- (a) Has its main effects on noradrenaline system in the brain
- (c) Works by affecting secondary messenger systems in the ^{glucagon} _{glucocorticoids}
- (d) Increases the rate of formation of cAMP
- (e) Works best with rapid cycling patients

81.

Side effects of lithium include the following except:

- (a) Leucocytosis
- (b) Acne
- (c) Alopecia
- (d) T wave inversion and QRS narrowing
- (e) Memory impairment

^{glucocorticoids}
via phosphatidyl inositol kinase 3.
synthase

Mut - tremor
Methimazole (potassium iodide)
D - lumbago
Ig problem - teratogenic

82.

Side effect of carbamazepine include all the following except:

- (a) Ataxia
- (b) Diplopia
- (c) SIADH (Chlom)
- (d) Agranulocytosis ✓
- (e) Leucocytosis

83.

The following are early signs (plasma levels 1.5 – 2 mEq/l) of lithium toxicity ^{Na+ GABA} except:

- (a) Anorexia ✓
- (b) Vomiting >
- (c) Diarrhoea ✓
- (d) Coarse tremor ✓
- (e) Nystagmus

84.

The following statements are true about carbamazepine ^{Na+ GABA} except:

- (a) Is a GABA agonist ✓
- (b) Affects calcium channels (dihydroperazine)
- (c) Induces its own metabolism ✓
- (d) Has a short half-life ✓ & valproic acid
- (e) Affects brain 5-HT function ✓

85.

Which statement is ^{Na+} SE concerning lamotrigine? ^{Na+}

- (a) Seems to be more effective in treating depressive episodes of bipolar ✓
- (b) Used less than other anticonvulsants for Bipolar Disorder
- (c) Voltage-gated sodium channel agonist
- (d) Inhibits the release of glutamate
- (e) Does not cause a rash as one of the main side effects

80. Which of the following statements is true concerning lithium?

- (a) Has its major effects on noradrenaline systems in the brain
- (c) Works by affecting secondary messenger systems in the cell via phosphatidylinositol d
- (d) Increases the rate of formation of cAMP glycogen synthase kinase 3.
- (e) Works best with rapid cycling patients

81. Side effects of lithium include the following except:

- (a) Leucocytosis
- (b) Acne
- (c) Alopecia
- (d) T wave inversion and QRS narrowing
- (e) Memory impairment

Mvt - tremor
Methotrexate (polyuria, polyuria)
O - Lupus like syndrome
Pq problem - telangiectatic

82. Side effect of carbamazepine include all the following except:

- (a) Ataxia
- (b) Diplopia
- (c) SIADH (水利尿)
- (d) Agranulocytosis ✓
- (e) Leucocytosis

83. The following are early signs (plasma levels 1.5 – 2 mEq/l) of lithium toxicity except:

- (a) Anorexia ✓
- (b) Vomiting ✓
- (c) Diarrhoea ✓
- (d) Coarse tremor ✓
- (e) Nystagmus

84. The following statements are true about carbamazepine except: $\text{Na}^+ \text{ GBT}$

- (a) Is a GABA agonist ✓
- (b) Affects calcium channels (dihydropyridines)
- (c) Induces its own metabolism ✓
- (d) Has a short half-life ✓ & valproic acid
- (e) Affects brain 5-HT function ✓

85. Which statement is FALSE concerning lamotrigine? Na^+

- (a) Seems to be more effective in treating depressive episodes of bipolar ✓
- (b) Used less than other anticonvulsants for Bipolar Disorder
- (c) Voltage-gated sodium channel agonist
- (d) Inhibits the release of glutamate
- (e) Does not cause a rash as one of the main side effects

86. Neuroleptic malignant syndrome is characterized by all of the following signs and symptoms except:

- (a) High fever ✓
- (b) Nausea and vomiting ✓
- (c) Automatic instability ✓
- (d) Muscle breakdown ✓
- (e) Elevated creatinine phosphokinase levels ✓

87. Risk factors for suicide may include all except:

- (a) Low income
- (b) Being an atheist
- (c) Family history of suicide
- (d) Chronic physical illness
- (e) Single and divorced marital status

88. Recognised roles of a physician does not include which ONE of these?

- (a) Promote well being
- (b) Prevent disease/disorder
- (c) To always provide custodial care for the sick
- (d) Enhance recovery
- (e) Promote human rights

89. Voluntary Kenya Community based organizations that are directly used as therapeutic agencies in mental health include the following except:

- (a) Alcoholic anonymous
- (b) Narcotic anonymous
- (c) Bipolar anonymous
- (d) Kenya association for the welfare of epileptics
- (e) Schizophrenic foundation

90. The national government has the following roles in organization of mental health services:

- (a) To improve government's responsiveness to local needs
- (b) Enhance effectiveness and efficiency of hospitals management
- (c) A B D E are all true
- (d) Increase the role of local community in management of hospitals/facilities
- (e) Reduce mortality, morbidity and disability for persons with psychological/physical disorders

91. Examples of psychiatric emergencies include the following except:

- (a) Violent aggressive patient
- (b) Last stages of remission
- (c) Suicidal patient
- (d) Patient in coma or semi coma
- (e) Psychosocial crisis

114. Delirium tremens, which can develop in persons who abstain from drinking a prolonged period of alcohol use, is characteristically associated with all of the EXCEPT:

- (a) Bradycardia
- (b) Tremor
- (c) Vivid visual hallucinations
- (d) Disorientation to time and place
- (e) A course of 3 to 7 days

* 115. True statements about the nature and effects of caffeine include all the following EXCEPT:

- (a) It often worsens the symptom occur with sudden cessation of chronic use
- (b) Withdrawal symptoms occur with sudden cessation of chronic use
- (c) Flashbacks occur with toxic reactions secondary to overuse
- (d) Overdose is associated with anxiety, derealization, delusions and tinnitus
- (e) Half-life of many caffeinated substance is about 3 to 7 h

116. Which of the following drugs is a tricyclic antidepressant?

- (a) Fluoxetine (Prozac)
- (b) Nortriptyline (Pamelor, Aventyl)
- (c) Phenelzine (Nardil)
- (d) Tranylcypromine (Parnate)
- (e) Clorazepate (Klonopin)

* 117. A 25 year old woman gives history of having used 30 mg/day of diazepam (Valium) for the last 20 months. Which of the following statement is most likely to be true?

- (a) There is a small chance she is physically dependant
- (b) She is almost certainly physically dependent
- (c) She is probably not physically dependant
- (d) She is probably not physically dependant, but psychologically habituated
- (e) Concern about physical dependency is not necessary at this dosage level

118. Side effects of antipsychotic drugs include all of the following EXCEPT:

- (a) Priapism
- (b) Galactorrhea
- (c) Amenorrhea
- (d) Retrograde ejaculation
- (e) Increased appetite and weight gain

119. All the following statements about electroconvulsive therapy (ECT) are true EXCEPT:

- (a) The principal indication is for the treatment of severe depression
- (b) It may be particularly effective in patients with delusional depression
- (c) It may be of benefit in the treatment of manic excitement
- (d) It is a procedure with a relatively high mortality
- (e) It was associated with impairment of memory

114

Delirium tremens, which can develop in persons who abstain from drinking after a prolonged period of alcohol use, is characteristically associated with all of the EXCEPT:

- a) Bradycardia \downarrow HR
- b) Tremor \checkmark
- c) Vivid visual hallucinations \checkmark
- d) Disorientation to time and place \checkmark
- e) A course of 3 to 7 days \checkmark

115

True statements about the nature and effects of caffeine include all the following EXCEPT:

- a) It often worsens ~~as~~ symptoms occur with sudden cessation of chronic use
- b) Withdrawal symptoms occur with sudden cessation of chronic use
- c) Flashbacks occur with toxic reactions secondary to ~~overdose~~
- d) Overdose is associated with anxiety, derealization, ~~disgust~~, and tinnitus
- e) Half-life of many caffeine-laden substances is about 3 to 7 h

116

Which of the following drugs is a tricyclic antidepressant?

- a) Fluoxetine (Prozac) ~~SSR~~
- b) Nortriptyline (Pamelor, Aventyl)
- c) Phenelzine (Nardil) ~~MAOI~~
- d) Tranylcypromine (Parnate)
- e) Clonazepam (Klonopin) ~~Benz~~

117

A 25 year old woman gives history of having used 30 mg/day of diazepam (\downarrow) over the last 20 months. Which of the following statement is most likely to be true?

- a) There is a small chance she is physically dependent
- b) She is almost certainly physically dependent
- c) She is probably not physically dependent
- d) She is probably not physically dependent, but psychologically habituated
- e) Concern about physical dependency is not necessary at this dosage level

118

Side effects of antipsychotic drugs include all of the following EXCEPT:

- a) Priapism
- b) Galactorrhea \checkmark
- c) Amenorrhea \checkmark
- d) Retrograde amnesia \checkmark
- e) Increased appetite and weight gain \checkmark

119

All the following statements about electroconvulsive therapy (ECT) are true EXCEPT:

- a) The principal indication is for the treatment of severe depression
- b) It may be particularly effective in patients with catatonic depression
- c) It may be of benefit in the treatment of manic excitement
- d) It is a procedure with a relatively high mortality \times
- e) It was associated with impairment of memory

86. Neuroleptic malignant syndrome is characterized by all of the following signs and symptoms except:

- (a) High fever ✓
- (b) Nausea and vomiting ✓
- (c) Automatic instability ✓
- (d) Muscle breakdown ✓
- (e) Elevated creatinine phosphokinase titres ✓

87. Risk factors for suicide may include all except:

- (a) Low income
- (b) Being an atheist
- (c) Family history of suicide
- (d) Chronic physical illness
- (e) Single and divorced marital status

T.D.S., T.R.E.
H.M.
Delusion

88. Recognised roles of a physician does not include which ONE of these?

- (a) Promote well being
- (b) Prevent disease/disorder
- (c) To always provide custodial care for the sick
- (d) Enhance recovery
- (e) Promote human rights

89. Voluntary Kenya Community based organizations that are directly used as therapeutic agencies in mental health include the following except:

- (a) Alcoholic anonymous
- (b) Narcotic anonymous
- (c) Bipolar anonymous
- (d) Kenya association for the welfare of epileptics
- (e) Schizophrenic foundation

90. The national government has the following roles in organization of mental health services:

- (a) To improve government's responsiveness to local needs
- (b) Enhance effectiveness and efficiency of hospitals management
- (c) A B D E are all true
- (d) Increase the role of local community in management of hospitals/facilities
- (e) Reduce mortality, morbidity and disability for persons with psychological/physical disorders

91. Examples of psychiatric emergencies include the following except:

- (a) Violent aggressive patient
- (b) Last stages of remission
- (c) Suicidal patient
- (d) Patient in coma or semi coma
- (e) Psychosocial crisis

120. Depression is not a uncommon side of:

- a) Insulin
- b) Cortisone
- c) Penicillin
- d) Imipramine (Tofranil)
- e) Bupropion (Wellbutrin)

*121. All of the following are symptoms commonly associated with tardive dyskinesia EXCEPT:

- a) Lip smacking or lip sucking
- b) Tongue movements
- c) Facial grimacing
- d) Fine tremors of the upper extremities
- e) Choreoathetoid movements of fingers and hands

122. All the following symptoms are associated with neuroleptic malignant syndrome EXCEPT:

- a) Hypothermia
- b) Rigidity
- c) Confusion
- d) Autonomic dysfunction
- e) Rhabdomyolysis

123. Side effects commonly associated with tricyclic antidepressants include all the following EXCEPT:

- a) Blurred vision
- b) Diarrhoea
- c) Dry mouth
- d) Urinary retention
- e) Tachycardia

124. Which of the following drugs is the least sedating?

- a) Chlorpromazine
- b) Imipramine
- c) Diazepam
- d) Lithium
- e) Haloperidol

125. The serum level of lithium at which therapeutic benefit levels off and side effects increase usually is considered to be:

- a) 0.3 Meq/l
- b) 1.0 Meq/l
- c) 1.5 Meq/l
- d) 2.0 Meq/l
- e) 3.0 Meq/l

0.5 - 1.0 mmol/L

126. Which of the following is not true about HIV/AIDS:

- a) People who suffers from schizophrenia have a higher risk of HIV infection compared to the general population
- b) Increased libido & sexual indiscretion in manic patients is predisposing factor
- c) Major depressive disorder does not necessarily predispose to HIV infection
- d) The neurotropic nature of the HIV virus increases the risk of psychiatric symptoms
- e) HIV positive persons are likely to use alcohol more often than the general population

127. Which is the most common form of stimulant medication for ADHD

- a) Vitalin
- b) Benzodiazepine
- c) Ritalin (methyl phenidate)
- d) Amylnitrate
- e) Epilium

128. Children with ADHD are known to have deficits in which of the following brain areas

- a) Perception
- b) Motor functioning
- c) Executive functioning
- d) Visual functioning
- e) Memory

129. Which of the following is correct about depression in children:

- a) Family therapy should be avoided because it scapegoats a child who is already vulnerable
- b) Symptoms may manifest as antisocial behaviour
- c) Anti-depressants generally are not effective in children
- d) The suicide rate in children 8 – 13 years is higher than it is in older adolescents
- e) Depression in children has been shown to be a prodrome to the later development of schizophrenia

* 130. Which of the following is behavior technique that has been successfully adapted to treat anxiety based problems in children:

- a) Systematic desensitization
- b) Systemic family therapy
- c) Electro convulsive therapy
- d) Physical therapy
- e) Online therapy

109. The following is true of alcohol intoxication and sexual performance except:

- (a) It impairs sexual performance
- ~~(b)~~ It causes poor erections
- (c) It decreases sexual desire
- (d) It reduces sperm motility
- (e) It increases sexual risk taking

110. Under the following condition, variation in sexual behaviour is considered pathological - Sexual Activity involving:-

- (a) Bonding in consenting adults
- (b) A 65 year old woman and a 22 year old boy
- ~~(c)~~ Consenting adult members of a nuclear family
- (d) A 65 year old man and a 22 year old boy
- (e) A consenting stranger

111. The following statements are true with regards to paraphilic except:

- (a) Most paraphilic disorders are very resistant to treatment ✓
- (b) Some are considered crimes in most countries ✓
- (c) Antidepressants particularly the selective serotonin reuptake inhibitors (SSRIs) ✓
- ~~(d)~~ Other forms of treatment include - ECT
- (e) Drugs that reduce testosterone levels in males may be used to treat some forms of paraphilic disorders ✓

112. A 45 year old married man has a telescope which he uses to watch women undressing in the opposite block at night. This behaviour is disrupting his marriage. The most likely diagnosis is:

- (a) Exhibitionism
- (b) Paedophilia
- (c) Impotence (erectile dysfunction)
- ~~(d)~~ Voyeurism
- (e) Sexual sadism

113. About 50% of all cases of sexual abuse develop PTSD. The main features of PTSD include the following except:

- (a) Re experiencing
- (b) Intrusive thoughts
- ~~(c)~~ Victimization
- (d) Avoidance
- (e) Hyperarousal

104. The following are known causes of sexual dysfunction in males except:
- (a) Diabetes mellitus
 - (b) Hypertension
 - (c) Antidepressants
 - (d) Vasectomy
 - (e) Antihypertensives

105. The following are known causes of dyspareunia in females except:
- (a) Poor arousal technique
 - (b) Inadequate vaginal secretions
 - (c) Use of drying herbs in the vagina
 - (d) Vaginitis
 - (e) Use of female condoms

Questions 106 - 108

Mary, aged 15 years has lost 10kg in 6 weeks. She is very thin but excessively concerned about being overweight. Her daily intake is 10 cups of coffee:

106. Mary might have the following symptoms except:
- (a) A reduced sex drive
 - (b) Increased sporting activities
 - (c) Amenorrhea
 - (d) May find herself very sexually attractive
 - (e) Increased body hair.

107. The most likely diagnosis would be:
- (a) Fetishism
 - (b) Body dysmorphic disorder
 - (c) Generalised anxiety disorder
 - (d) Bulimia Nervosa
 - (e) Anorexia Nervosa

108. The following have been suggested in the etiology of the above disorder except:
- (a) Hereditary factors
 - (b) Socio cultural factors
 - (c) History of sexual abuse
 - (d) Intellectual factors
 - (e) Racial factors

119. Sub cortical dementias include the following except:

- (a) Parkinson's disease
- (b) Huntington's disease
- (c) Wilson's disease
- (d) Pick's disease
- (e) Binswanger's disease

120. The commonest cause of dementia is:

- (a) Dementia with Lewy body
- (b) Frontal temporal dementia
- (c) Alzheimer's dementia
- (d) Vascular dementia
- (e) Creutzfeldt - Jakob disease

121. Predisposing factors to Alzheimer's disease include the following except:

- (a) Male sex
- (b) Head injury
- (c) Down's syndrome
- (d) Positive family history
- (e) Post menopausal estrogen decline

122. Clinical features of Lewy body dementia include the following except:

- (a) Prominent auditory hallucinations
- (b) Vivid visual hallucinations
- (c) Delirium
- (d) Parkinsonism features
- (e) Disturbed rapid eye movement (REM)

123. Potentially reversible causes of dementia include the following except:

- (a) Normal pressure hydrocephalus
- (b) General paralysis of the insane
- (c) Myxoedema
- (d) Semantic dementia - semantic - old
- (e) Subdural hematoma

124. Combinations of antipsychotic drugs:

- (a) Should never be used
- (b) Can be used in special cases
- (c) Are invariably toxic
- (d) Increase the risk of relapse directly
- (e) Should be used to treat recent onset or newly diagnosed schizophrenia

114. The following are specific strategies that are commonly used during psychosexual therapy except:

- (a) Sensate Couple focus
- (b) Exploration of different coitus positions
- (c) Use of lubricants
- (d) Semaan's technique
- (e) Exchanging partners

115. NOT TRUE: Perpetrators of sexual abuse are usually:

- (a) Mentally ill
- (b) Plan and plot overtime if known to child
- (c) Usually target children who are shy and have low self esteem
- (d) Males close to and known to the children
- (e) Are represented in all socio economic groups

116. Which one of the following is correctly matched in terms of degree of risk of transmission of HIV:

- (a) Anal sex – Moderate risk
- (b) Vaginal sex – Higher risk than anal sex
- (c) Oral sex – Moderate risk
- (d) Scarification – No risk
- (e) Sharing needles – High risk

117. Which of the following is not true of cultural practices that have been implicated in the transmission of HIV?

- (a) Female genital mutilation
- (b) Polygamous marriages
- (c) Marriage of adolescent girls by older men
- (d) Sharing ceremonial beer from one drinking utensil
- (e) Wife inheritance

118. Social interventions for the management of an episode of schizophrenia include all of the following except:

- (a) Addressing need for time off work/school
- (b) Protection of the patients social support networks
- (c) Providing public education that decrease/prevent mental illness related stigma
- (d) Improving social networks
- Limit contact with family during the whole period of time that a patient is symptomatic



4
5
4
3

UNIVERSITY OF NAIROBI

UNIVERSITY EXAMINATIONS 2017/2018

YEAR 5 EXAMINATIONS FOR THE DEGREE OF BACHELOR OF MEDICINE AND BACHELOR OF SURGERY

MCQS - PSYCHIATRY

DATE: MARCH 26, 2018

TIME: 9.00 A.M. - 11.00 A.M.

INSTRUCTIONS

- 1) Each question contains 5 suggested responses. Select the best response to each question.
- 2) Use the answer sheet provided to circle the correct response.
- 3) There is no penalty mark for wrong answer.

1. Which ONE of the following statements is the best example of a good interview skill?

- (a) Asking closed questions during the interview
- (b) Being sympathetic rather than empathetic
- (c) Maintaining continuous eye contact
- (d) Premature assurances which are helpful
- (e) The use of non-verbal methods of communication ✓

2. Which of the following stages of Freud's theory of psychosexual development already characterised Erik Erikson's Initiative versus Guilt stage?

- (a) Anal (2-3 yrs)
- (b) Genital (Puberty to adulthood)
- (c) Latency (age 6 to puberty)
- (d) Oral (0-12 months)
- (e) Phallic (3-5 yrs)



① Infancy (1-2 yrs)

3. The following are considered founders of modern psychiatry except:

- (a) Ivan Pavlov ✓
- (b) Jean Larivee Dominique Esquirol ✓
- (c) Philippe Pinel ✓
- (d) Paul Kraepelin ✓
- (e) Eugene Bleuler ✓

Page 1 of 29

Adolescence



Erikson classification

- ① Infancy (0-18 mos) - Basic Trust vs. Mistrust (0-18 mos)
- ② Autonomy vs. shared doubt (1-3 yrs)
- ③ Initiative vs. guilt → (pre-school) (3-5 yrs)
- ④ Industry vs. Inferiority - (5-12 yrs) - Hardworking vs. Laziness
- ⑤ Identity vs. Role confusion - (12-18 yrs) - Adolescence
- ⑥ Intimacy vs. Isolation - (18-30 yrs)
- ⑦ Generativity vs. Stagnation - (30-65 yrs)
- ⑧ Ego Integrity vs. Despair - (65 and above)

4. The following are associated with behaviour therapy except:

- (a) B.F Skinner - Theory of behaviorism
(b) Joseph Wolpe - Behavior therapy
~~(c)~~ Jean Piaget - cognitive development
(d) J.B Watson - Psychological school of behaviorism
~~(e)~~ A. Lazarus - cognitive therapy

5. The assertion that all behaviours are driven by antecedent events, experiences.
There are no accidents; nothing happens by chance is known as:

- ~~(a)~~ Psychic determinism
(b) Learned helplessness
(c) Unconditional positive regard
(d) Free association
(e) Unconscious

6. What is transference?

- (a) Displacement of feelings from future experiences to past experiences
(b) Displacement of ideas from mother to child
~~(c)~~ Displacement of feelings, attitudes, behavioural expectations and attributes from important childhood relationships to current ones
(d) Transferring material resources to a client's bank account
(e) Defense mechanism that helps to keep tab on feelings of security

7. Counter transference refers to:

- ~~(a)~~ The therapist projecting their own feelings ("issues," emotional baggage, etc.) onto their patient
The patient projecting their own feelings (issues, emotional baggage) onto the therapist
(c) The child projecting his/her feelings on the parents and parental rejection of those feelings
Considering the transference made by the patient through a systematic analysis
(e) Displacement of feelings, attitudes, behavioural expectations and attributes from important childhood relationships to current ones

8. Unconsciously assisting others to avoid negative personal feelings refers to:

- ~~(a)~~ Sublimation
~~(b)~~ Altruism
(c) Displacement
(d) Condensation
(e) Projection



9.

Which one of these techniques is the fundamental base of any psychotherapy practice?

- (a) Psychoanalyzing
- (b) Dream analysis
- (c) Empathy
- (d) Punishment
- (e) Systemic desensitization

10.

Which one of the following techniques is associated with behavioral psychotherapy?

- (a) Fertilization
- (b) Facial coding
- (c) Fabrication
- (d) Flooding
- (e) Psychoeducation

11.

One of the client centered ideas given by Carl Rogers includes:

- (a) Unconscious
- (b) Preconscious
- (c) Existentialism
- (d) Unconditional Positive Regard
- (e) Conditional Negative Regard

12.

In selecting a client to join a group therapy, the following can lead to exclusion from the group except:

- (a) Unwillingness or low motivation to participate
- (b) Extremely elevated level of distress
- (c) Member will benefit from the group experience
- (d) Member will not be able to listen to other members
- (e) Non compliance with group rules

13.

Which ONE is true about pregnancy?

Pregnancy affects the psychological well being of all women

(b) Pregnancy affects the psychological well-being of only those women not prepared for the pregnancy

(c) The psychological well-being during pregnancy is an only woman's affair

(d) The psychological concerns of the pregnant woman do not need to be addressed at all

(e) None of the above

- 24 Which ONE of the following is NOT a general aim in CBT? *cognitive behavioral flora*
- (a) Modifying maladaptive behaviour patterns
(b) Eliminating completely environmental conditions that may be causing or maintaining such problematic behaviors
(c) Improving clients' skills related to work and social interactions
(d) Resolve some inner conflicts and stress handicapping and disabling the client
(e) Restructuring a person's distorted cognition
- 25 Which ONE of the following statements is true about CBT?
- (a) All mentally ill persons can be managed using CBT approach
(b) CBT process starts with behaviour modifications before restructuring client's distorted belief system
(c) Some of the more responsive subjects one may not remember the process
(d) Clients recruited for CBT always report feeling bored
(e) CBT is applied among clients with alcohol problems so that social skills are built on concrete process
- 26 The following CBT techniques are applied during CBT except:
- (a) Listening as the patients talk about their emotional difficulties
(b) Assertive training
(c) Systematic desensitization
(d) Contingency management
(e) Token economy
- 27 Which of the following statements is true about aims of family therapy?
- (a) Therapist induces Arguments
(b) Therapy improves communication
(c) Therapy decreases autonomy for each member
(d) The therapy does not improve agreement about roles
(e) The therapy increases aggression in the member who is the patient
- 28 Which one of the following is not true about models for family therapy?
- (a)* One member is treated individually by another therapist
(b) Group therapy is carried out in a family therapy session
(c) Couples learn from listening to other families' members
(d) Family members are seen together
(e) Individual therapy can go on with the same therapist but at different times.
- 29 Family therapy is not indicated in following situations
- (a) Communication problems between family members
(b) Problem by the child
(c) Treatment of young persons with Anorexia Nervosa
(d) Relapse prevention in Schizophrenia
(e) Manic phase of a member with Bipolar I mood disorder

30. The following roles is not played by a family therapist:
- (a) Identifying the family disagreement
 - (b) Helping a family to understand and modify the rules
 - ~~(c)~~ Identifying age of children
 - (d) Identify the families with distorted ways of communicating
 - (e) Identifying the family unspoken rules
31. The approach during family therapy is that:
- (a) Therapist focuses on one family member
 - (b) Therapist does not encourage family members to speak to each other
 - (c) System tasks focus on behaviours which occur rarely
 - (d) Family members are not involved in organizing a timetable indicating when this would take place, duration of it and frequency
 - ~~(e)~~ Therapist must set up tasks and injunctions
32. Emil Kraepelin classified the following disorders except:
- (a) Dementia praecox
 - (b) Catatonic
 - (c) Hiebephrenic
 - (d) Paranoid
 - ~~(e)~~ Cti du chat syndrome
33. Your patient tells you that he has suddenly remembered being taken into hospital as a child and having a microchip implanted in his brain. Of what is this example?
- ~~(a)~~ Autochthonous delusion
 - (b) Delusion of reference
 - (c) Delusion memory
 - (d) Delusion mood
 - (e) Delusion percept
34. Which ONE of the following is a 'neurological soft sign'?
- ~~(a)~~ Cognitive impairment
 - (b) Dysbindin genotype
 - (c) Impaired fist-edge-palm test
 - (d) Positive stroop test
 - ~~(e)~~ Receptive dysphasia
35. The following are psychological treatments of depression except:
- (a) Cognitive behaviour therapy
 - (b) Group therapy
 - (c) Individual psychotherapy
 - (d) Family therapy
 - ~~(e)~~ Electroconvulsive therapy

14. The following are correct about group therapy except:

- (a) The group therapy must be run by a professionally trained mental health worker
- (b) Group therapy is the same as support group
- (c) Group therapy is effective in treatment of mental disorders
- (d) Group therapy must be structured
- (e) Group therapy is based on theoretical approach depending on the mental health problem

15. Exposure therapy is an effective behavioural management technique that is useful in managing all of the following mental health problems in CBT (Cognitive Behaviour Therapy) except:

- (a) Phobias
- (b) Relationship difficulties
- (c) Schizophrenia
- (d) Sleeping disorders
- (e) Eating disorders

16. A number of aspects of learning are used in the principles of CBT (Cognitive Behaviour Therapy). Which of the following is NOT a principle of learning during CBT?

- (a) Operant conditioning
- (b) Classical conditioning
- (c) Negative Reinforcement
- (d) Social learning

17. Relaxation is shown to be effective in managing anxiety. Which of the following is NOT a relaxation exercise that has been researched on in connection with CBT (Cognitive Behaviour Therapy)?

- (a) Progressive muscle relaxation
- (b) Progressive Anxiety relaxation
- (c) Yoga
- (d) Diaphragmatic breathing relaxation
- (e) Imagery

18. According to CBT (Cognitive Behaviour Therapy) where do the early maladaptive schemas or bottom lines originate from?

- (a) Current negative experiences and relationships
- (b) Early childhood negative experiences and relationships
- (c) Peer pressure and friendship influences
- (d) The cognitive behaviour therapy done in the sessions
- (e) In early adulthood years as an adult starts managing their livelihoods

19. Behaviour therapy in CBT (Cognitive Behaviour Therapy) is based on all of the following except:

- (a) Operant conditioning
- (b) Scientific knowledge of human nature
- (c) Classical conditioning
- (d) Only on Tradition western traditions
- (e) Learning principles

20. Behaviour therapy in CBT focuses on all of the following except:

- (a) Self awareness
- (b) Eliminating symptom
- (c) Learning adaptive behavioural skills
- (d) Searches of conscious conflicts
- (e) Self evaluation

21. Seeking to understand the message of a patient by asking for more information or for elaboration on a point during CBT process is referred to as?

- (a) Social response
- (b) Minimal cue
- (c) False reassurance
- (d) Clarification
- (e) Amplification

22. Behavioral aspect in Cognitive Behavioral Therapy is not used to facilitate change in:

- (a) Thought stopping
- (b) Relaxation
- (c) Symbolisation
- (d) Repeated irrational thoughts
- (e) Examining thoughts of the client during therapy

23. One of the following psychological problems cannot be managed using CBT. Which ONE is it?

- (a) Eating disorder
- (b) Depression
- (c) SADS (Seasonal Affective Disorder)
- (d) PTSD
- (e) Dementia

35. A woman who has lost a pregnancy can experience all of the following except:
- (a) Guilt
 - (b) Personal failure
 - (c) Inadequacy
 - (d) Positivity
 - (e) Anger

37. Prolonged grieving due to loss of pregnancy can occur due to:
- (a) The loss and grief being minimized by those around her
 - (b) Seeing or being allowed to see the foetus or baby
 - (c) Getting support from the partner
 - (d) Knowing the cause of the loss of pregnancy
 - (e) Not worried about future pregnancies

38. The most widely prescribed pharmacological treatment for attention deficit hyperactivity disorder (ADHD) is:
- (a) Diazepam
 - (b) Ritalin
 - (c) Lithium
 - (d) Risperidone
 - (e) Largactil

39. Altruism, humour and sublimation are part of which of the following defence mechanisms?
- (a) Immature
 - (b) Mature
 - (c) Neurotic
 - (d) Narcissistic
 - (e) Psychotic

40. Patients with psychogenic pain disorder have the following characteristics except.
- (a) Have ongoing pain, not due neoplastic disease
 - (b) Have no significant ongoing pathophysiological mechanisms that account for the ongoing pain
 - (c) Have a life history of inability to form any psychological view of life problems
 - (d) Their chronic pain has become the central focus to their thoughts, feelings, behavior and social relationships
 - (e) They are not amenable to cognitive behavioral theory

41 All of the following are specific somatoform disorders listed in DSMIV TR except:

- (a) Conversion disorder ✓
- (b) Pain disorder
- (c) Hypochondriasis ✓
- (d) Body dysmorphic disorder
- ~~(e)~~ Differentiated somatoform disorder

42 A 30 year old female prisoner when invited by the prison medical officer to come into the interview room takes four steps backwards from the door. She came into the room when the officer asked her to go away. Which ONE of the following behavioural disorders of schizophrenia is she exhibiting?

- (a) Advertence
- ~~(b)~~ Ambitendency
- ~~(c)~~ Negativism
- (d) Satyriasis
- (e) None of the above

43 Which ONE is odd amongst the following?

- (a) Paroxetine
- (b) Fluoxetine
- (c) Fluvoxamine
- (d) Sertraline
- ~~(e)~~ Rispexone

44 A 40 year old female with a long history of Schizophrenia relapses. On examination she holds her arm and face in a peculiar posture which is maintained for a few minutes at a time. What motor disorder is she exhibiting?

- (a) Ambitendency
- ~~(b)~~ Mannerism
- (c) Chorea
- (d) Stereotypy
- ~~(e)~~ None of the above

Posturing

45 A 26 year old man is diagnosed with Schizophrenia. Which one of the following predicts a poor prognosis?

- (a) Absence of negative symptoms
- (b) Being married
- (c) Few or no episodes in the past
- ~~(d)~~ Good premorbid personality
- ~~(e)~~ Gradual onset

For questions 46 - 48

A 7 year old boy is brought to you by his parents on a referral by the school where the child is in class one. The boy does not have a major discipline problem but he frequently answers questions without being called on and is often out of his seat without permission. His school work is adequate but the teacher believes, "he could do better." He has difficulty completing tasks and appears to spend most of the class time day dreaming.

46. Which additional piece of information would support the most likely etiology for his symptoms?

- (a) A history of head injuries
- (b) A history of neurological symptoms
- (c) A history of tics
- (d) His medication history
- (e) Family psychiatry history

47. The most likely diagnosis is:

- (a) Attention deficit hyperactivity disorder (ADHD)
- (b) Conduct disorder
- (c) Post Traumatic Stress Disorder
- (d) Intellectual disability
- (e) Autism spectrum disorder

48. Regarding treatment the best advice to the family would be that:

- (a) He has a diagnosable disorder so he should not be held accountable for his symptoms
- (b) He should alter his diet immediately
- (c) He needs intensive, probably long term psychotherapy
- (d) Medication might be helpful
- (e) They should probably NOT discuss his diagnosis with the teacher as it might be stigmatizing

49. Which of the following is among the most common projective techniques used in a child mental health interview in order for the child to express concerns indirectly?

- (a) Asking questions about specific symptoms
- (b) Drawing
- (c) Have the child climb stairs to assess gross motor development
- (d) Asking the child how he feels
- (e) Assessing fund of knowledge

50. The psychiatric assessment of the child commonly differs from the assessment of adults in all of the following ways except:
- (a) The child and the clinician are at different developmental levels such that they may have difficulties in communication
 - (b) The clinician need only focus on the assessment and treatment of the child
 - (c) The child may function differently at different settings
 - (d) The child's presenting problems must be examined in a developmental context
 - (e) There are developmental differences in the presentation of mental illness as categorized in DSM-IV TR / DSM 5
51. Which of the following is considered the single strongest predictor of adolescent substance abuse?
- (a) Parental permissiveness
 - (b) Having friends who use drugs
 - (c) Low socio economic status
 - (d) History of poor academic performance
 - (e) Parental attitudes towards drinking and their drinking habits
52. Symbolic functioning, structure, content and expressions of and control of aggression are characteristics of which component of the population -- appropriate mental status exam for toddlers and infants?
- (a) Self-regulation
 - (b) Intellectual functioning
 - (c) Play
 - (d) Affect and mood
 - (e) Relatedness
53. An advantage of having a set of toys available in the office versus having the child bring toys from home is:
- (a) Observation frustration with toys too difficult for the child's developmental level
 - (b) The child will be uninterested in new toys and more engaged with interview
 - (c) The child will feel more at ease if the child is playing
 - (d) The child will have to adapt to the evaluator
 - (e) Limited distractions from the child's favourite toys
54. Which of the following is associated with better outcomes in childhood-onset Schizophrenia?
- (a) Early age of onset
 - (b) Non-acute insidious onset
 - (c) Presence of affective symptoms
 - (d) Low IQ
 - (e) Long duration of untreated psychosis

55. In people with intellectual disability, psychiatric disorders:

- (a) Are often over diagnosed
- (b) Diagnostic overshadowing completely explains the difference in prevalence
- (c) Diagnostic overshadowing is not due to bias of clinician
- (d) Diagnostic overshadowing is deliberate
- (e) Are often under diagnosed

56. Which ONE of the following is true regarding childhood schizophrenia?

- (a) Low socioeconomic status has been well documented to be a risk factor for childhood onset Schizophrenia
- (b) Bipolar disorder in children is frequently misdiagnosed as having Schizophrenia
- (c) Abuse or neglect does not affect one's risk for developing psychosis
- (d) Children with Autism Spectrum disorder possess pervasive hallucinations, delusions, and thought disorder characteristics similar to Schizophrenia
- (e) Hallucinations in autism are typically chronic

57. Which of the following would NOT be considered to be an essential part of a neuropsychological assessment of child?

- (a) Medical history
- (b) Social history
- (c) Behavioral observations of the child
- (d) Psychological testing
- (e) Magnetic Resonance Imaging (MRI)

58. A 7 year old girl is referred for an assessment due to worsening performance at school. She was recently diagnosed with epilepsy and her medications have required frequent adjustments to keep seizures under control. She is also noted to have become more socially withdrawn and spends less time with friends. What is the most likely contributor to her recent poor school performance?

- (a) Brain tumour
- (b) Depression
- (c) Medication side effect
- (d) Hydrocephalus
- (e) Substance abuse

55 With regards to the parents of the child being evaluated, a child psychiatrist should take all of the following considerations except:

- (a) The expectations as well as feelings about the evaluation should be elicited from the child's parents
- (b) Parents may bring a mixed heritage of guilt, shame, concern and fear of being judged in seeking help for their child
- (c) Parents may have unrealistically optimistic fantasies of "absolution" of unconscious guilt or hopes of quick cures
- (d) Parents generally inform the child of the nature of the evaluation prior to arrival
- (e) Parents generally have a well-formed understanding of psychiatric assessment and treatment from mass media sources

60. In which of the following situations would the risk of developing Schizophrenia be the lowest for a child?

- (a) A child's Schizophrenic parent 12% has a non Schizophrenic identical twin 0%
- (b) A child's non Schizophrenic parent 0% has a Schizophrenic identical twin 4%
- (c) A child's Schizophrenic parent 12% has a non Schizophrenic fraternal twin 0%
- (d) A child's non Schizophrenic parent 0% has a Schizophrenic fraternal twin 4%
- (e) A child's non-Schizophrenic parent 0% has a non Schizophrenic sibling 0%

61. The following are correct about children below the age of 7 years in relation to understanding death in the family except:

- (a) Cannot comprehend the irreversibility of death
- (b) Can express their feelings in play therapy
- (c) Has limited ability to verbalize their feelings
- The children tend towards magical thinking and ego centricity
- The children are sensitive about being 'different' from their peers

62. In attention deficit hyperactivity disorder (ADHD), which of the following is NOT classified as a problem relating to poor attention?

- (a) Appears routinely forgetful
- Experiences difficulties in waiting in turn
- Dislikes tasks involving mental effort e.g. homework
- Easily distracted
- Appears not to listen when spoken to

63. The following are depot antipsychotic preparations except:

- (a) Fluphenazine decanoate
- (b) Haloperidol decanoate
- (c) Olanzapine pamoate
- (d) Zuclopentixol decanoate
- (e) Clozapine decanoate

64. Regression of milestones is a common presentation of childhood disorders. Which of the following medical conditions is associated with regression? -

- (a) Central Nervous System infections
- (b) Phenylketonuria
- (c) Hypothyroidism
- (d) Seizure disorder
- (e) All of the above

65. HIV/AIDS is a risk factor for all of the following disorders except:

- (a) Schizophrenia
- (b) Alcohol use disorder
- (c) Depression
- (d) Adjustment disorder
- (e) Anxiety disorder

66. In addition to clinical and family history, a near confirmatory diagnostic test for tic disorders is:

- (a) Magnetic Resonance Imaging
- (b) Single photon emission computed tomography
- (c) Electroencephalogram
- (d) Blood dopamine level
- (e) No test

67. In the Four P's Model of psychiatric formulation, which domain of factors is concerned with features that make the presenting condition endure, such as the severity of the condition or compliance issues?

- (a) Predisposing
- (b) Precipitating
- (c) Perpetuating
- (d) Protective
- (e) Persistence

68. What is the strongest risk factor for youth suicide?

- (a) Major depression
- (b) Bipolar disorder
- (c) Substance use disorder
- (d) Prior history of a suicide attempt
- (e) Borderline personality disorder

69. What is a simple alternative to the bell and pad method for the treatment of enuresis?
- (a) Waking up the child using a bull horn
(b) Setting an alarm clock after 2 to 3 hours of sleep when the bladder may be reaching full capacity
(c) Keeping the lights on at night to aid arousal when the bladder signals full capacity
(d) Restricting fluids after 4 pm
(e) None of the above
70. The psychiatrist assessment of children commonly differs from the assessment of adults in ALL of the following ways except:
- (a) The child's behaviour may cause greater distress to the adults than to the child
(b) Children may not recognize their behaviour as problematic to others
(c) Children often seek out psychiatric assessment or treatment for themselves
Children may attribute problems to others and be unwilling to accept their own personal contribution to the problem
(e) The adult's expectation for the child may exceed the child's abilities
71. Which of the following symptoms of ADHD (Attention Deficit Hyperactivity Disorder) is most likely to be outgrown when a child matures to adulthood?
- (a) Inattention
(b) Hyperactivity
(c) Impulsivity
(d) Distractibility
(e) Both hyperactivity and impulsivity
72. A 74 year old man presents with memory impairment and behaviour that is out of character. Which of the following cognitive tests is appropriate to test frontal lobe function?
- (a) Benton Visual retention test
(b) National adult reading test
(c) Rey-Osterrieth Complex figure test
(d) Rorschach ink blot test
(e) Wisconsin Card Sorting test
73. A 76 year old woman suffers a stroke due to a ruptured berry aneurysm in the anterior cerebral artery and the frontal lobe is affected. Which ONE of the following tests would you expect to be abnormal?
- (c) Colour vision testing
(d) Copying intersecting pentagons
(e) Luria's motor test
(f) Semantic memory
(g) Three object recall

74. Which ONE of the following was the first effective antipsychotic therapy for Schizophrenia or bipolar disorder?

- (a) Chlorpromazine
~~(b)~~ ECT
(c) Hypnotherapy
(d) Insulin coma therapy
(e) Pyrotherapy

75. Which ONE of the following anti-depressants is NOT suitable for a patient who is suffering from depression and has recently also had myocardial infarction?

- ~~(A)~~ Amitriptyline
(b) Duloxetine
(c) Mirtazapine
(d) Paroxetine
(e) Sertraline

76. Which of the following Neurotransmitters promote 'Sleep'?

- (a) Acetylcholine
(b) Dopamine
(c) Histamine
(d) Noradrenaline
~~(e)~~ Serotonin

77. Which ONE of the following can be used to reduce alcohol consumption in patients who are still consuming alcohol?

- (a) Acamprosate
~~(b)~~ Bupropion
~~(c)~~ Disulfiram
(d) Nalmefene
(e) Naltrexone

78. In the use of antidepressants in the management of Bipolar disorder, which of the following induces a metabolic syndrome?

- (a) Lamotrigine
(b) Lithium
(c) Lorazepam
~~(d)~~ Olanzapine
(e) Sibutramine

79. In bipolar disorder, which of the following is useful in acute mania?

- ~~(A)~~ Bupropion
Carbamazepine
(c) Imipramine
(d) Sertraline
(e) Tricyclics

Which of the following statements is true concerning lithium?

- (a) Has its main effects on noradrenaline systems in the brain
- (b) Works by affecting secondary messenger systems in the cell
- (c) Increases the rate of formation of cAMP
- (d) Works best with rapid cycling patients
- (e) Is not absorbed from the kidney

81 Side effects of lithium include the following except:

- (a) Leucocytosis ✓
- (b) Acne ✓
- (c) Anorexia
- (d) T wave inversion and QRS narrowing
- (e) Memory impairment

82 Side effect of carbamazepine include all the following except:

- (a) Ataxia ✓
- (b) Diplopia
- (c) SIADH
- (d) Agranulocytosis✓
- (e) Leucocytosis

83 The following are early signs (plasma levels 1.5 – 2 mEq/l) of lithium toxicity except:

- (a) Anorexia ✓
- (b) Vomiting ✓
- (c) Diarrhoea
- (d) Coarse tremor ✓
- (e) Nystagmus

84. The following statements are true about carbamazepine except:

- (a) Is a GABA agonist
- (b) Affects calcium channels
- (c) Induces its own metabolism
- (d) Has a short half life
- (e) Affects brain 5-HT function

85. Which statement is FALSE concerning lamotrigine?

- (a) Seems to be more effective in treating depressive episodes of bipolar disorder
- (b) Used less than other anticonvulsants for Bipolar Disorder
- (c) Voltage-gated sodium channel agonist
- (d) Inhibits the release of glutamate
- (e) Does not cause a rash as one of the main side effects

86. Neuroleptic malignant syndrome is characterized by all of the following signs and symptoms except:

- (a) High fever
~~(b)~~ Nausea and vomiting
~~(c)~~ Automatic instability
(d) Muscle breakdown
(e) Elevated creatinine phosphokinase titres

87. Risk factors for suicide may include all except:

- (a) Low income
~~(b)~~ Being an atheist
(c) Family history of suicide
(d) Chronic physical illness
(e) Single and divorced marital status

88. Recognized roles of a physician does not include which ONE of these?

- (a) Promote well being
(b) Prevent disease/disorder
~~(c)~~ To always provide custodial care for the sick
(d) Facilitate recovery
(e) Promote human rights

89. Voluntary Kenya Community based organizations that are directly used as therapeutic agencies in mental health include the following except:

- (a) Alcoholic anonymous
(b) Narcotic anonymous
~~(c)~~ Bipolar anonymous
(d) Kenya association for the welfare of epileptics
(e) Schizophrenic foundation

90. The national government has the following roles in organization of mental health services:

- (a) To improve government's responsiveness to local needs
(b) Enhance effectiveness and efficiency of hospitals management
(c) A B D E are all true
(d) Increase the role of local community in management of hospitals/facilities
(e) Reduce mortality, morbidity and disability for persons with psychological/physical disorders

91. Examples of psychiatric emergencies include the following except:

- (a) Uncontrolled aggressive patient
(b) Late stages of remission
(c) Suicidal patient
(d) Patient in coma or semi-coma
(e) Psychosocial crisis

92. The following are illness causing behaviour except:

- (a) Cigarette smoking
- (b) Obesity
- (c) Promiscuity
- ~~(d)~~ Physical exercise
- (e) High fat diet

93. The capacity to form concepts and generalize items is called:

- ~~(a)~~ Concrete thinking
- ~~(b)~~ Abstract thinking
- (c) Delusional thinking
- (d) Intellectualization
- (e) Rationalization

94. Psychological disorders are influenced by the following except:

- (a) Learned helplessness
- (b) Trauma
- (c) Role problems
- ~~(d)~~ Stress
- ~~(e)~~ Cohesiveness

95. Children with ADHD are known to have deficits in which of the following brain areas:

- (a) Perception
- (b) Motor functioning
- ~~(c)~~ Attention
- (d) Memory
- (e) Sensory functioning

96. Which of the following perinatal risk factors are not associated with attention deficit hyperactivity Disorder (ADHD)?

- (a) Low birth weight
- ~~(b)~~ Maternal alcohol misuse ✓
- ~~(c)~~ Measles, mumps and rubella vaccines
- (d) Parents who smoke during pregnancy increase the risk of having a child with ADHD
- (e) Maternal stress

97. The following are signs of alcohol dependence except:

- ~~(a)~~ Tolerance
- ~~(b)~~ Withdrawal symptoms
- ~~(c)~~ Hypersomnia
- (d) Change of priorities
- (e) Reinstatement to dependence after a period of abstinence

119. Sub cortical dementias include the following except:

- (a) Parkinson's disease
- (b) Huntington's disease
- (c) Wilson's disease
- (d) ~~Pick's disease~~ — frontotemporal
- (e) Binswanger's disease

120. The commonest cause of dementia is:

- (a) Dementia with Lewy body
- (b) Frontal temporal dementia →
- (c) ~~Alzheimer's dementia~~
- (d) Vascular dementia
- (e) Creutzfeldt - Jacob disease

121. Predisposing factors to Alzheimer's disease include the following except:

- (a) Male sex ✓
- (b) Head injury ✓
- (c) Down's syndrome ✓
- (d) Positive family history ✓
- (e) Post menopausal estrogen decline ✓ → HRT

122. Clinical features of Lewy body dementia include the following except:

- (a) Precipitant auditory hallucinations
- (b) Vivid visual hallucinations ✓
- (c) Delirium ✓
- (d) Parkinsonism features ✓
- (e) Disturbed rapid eye movement (REM) ✓

123. Potentially reversible causes of dementia include the following except:

- (a) Normal pressure hydrocephalus ✓
- (b) Glioma/pituitary adenoma
- (c) Myxoedema
- (d) Semantic dementia
- (e) Subdural hematoma ✓

124. Combinations of antipsychotic drugs:

- (a) Should never be used
- (b) Can be used in special cases
- (c) At a low dose
- (d) Increase the risk of relapse directly
- (e) Should be used to treat, recent onset or newly diagnosed schizophrenia

125. Regarding the treatment of anxiety disorders:

- (a) Antipsychotic drugs remain the first line choice
- (b) Antidepressants when used correctly, offer rapid relief
- (c) Benzodiazepines should never be used because of their addictive properties
- (d) The dose of tricyclic antidepressants should be reduced gradually to avoid rebound effects
- (e) An apparent worsening of symptoms may be seen when starting treatment with anti depressants

126. Useful interview techniques include: except

- (a) Summation
- (b) Silence
- (c) Missing appointments
- (d) Homework
- (e) Directive questioning

127. In reviewing a psychotic patient the following are important: *Choose the best response.*

- (a) Medications used, their doses and side effects
- (b) Social needs
- (c) The history from the caretaker
- (d) Suicidal tendencies
- (e) Physical health

128. The importance of an objective history is demonstrated in establishing the diagnosis of sleep apnea. The patient's bed partner, though not necessarily the patient, is likely to report all the following except.

- (a) Agitated behaviour
- (b) Loud snoring
- (c) Sleep walking
- (d) Gasping
- (e) Bed wetting

129. Regarding a person with alcohol abuse of the following are true except

- (a) Recurrent legal problems related to alcohol
- (b) Continued use despite recurrent interpersonal or social problems
- (c) Failure to fulfill role obligation at work or home because of excessive drinking
- (d) Stop to abuse alcohol if advised to stop
- (e) Accidents are common

130. All of the following are symptoms of depression except

- (a) Anhedonia
- (b) Terminal insomnia
- (c) Thought broadcasting
- (d) Loss of interest to all pleasurable activities
- (e) Nihilistic delusions

131. 'Maternity blues' is accurately characterized by which of the following?

- (a) It is more acute than post partum depression
- (b) It is usually a chronic and relapsing syndrome
- (c) It affects 50 – 80% of all new mothers
- (d) It is characterised by persistent apathy
- (e) It is not associated with sleep disturbance

132. The basis for the therapeutic effect of electroconvulsive therapy (ECT) is:

- (a) Seizure activity
- (b) Electrical stimulation of the brain
- (c) Memory loss
- (d) The depressed patient's wish for punishment
- (e) The depressed patient's attitude toward ECT

133. The following investigation is ~~not~~ mandatory before electroconvulsive therapy:

- (a) EEG
- (b) Liver function test
- (c) Urea and electrolytes
- (d) Urinalysis
- (e) Fasting blood sugar

134. Electroconvulsive therapy is contraindicated in:

- (a) Pregnancy
- (b) Persons with recent myocardial infarction
- (c) Persons with controlled hypertension
- (d) Persons with history of febrile convulsions in childhood
- (e) Persons aged above 75 years

135. The following is a parasomnia:

- (a) Nighttime disorder
- (b) Narcolepsy
- (c) Primary insomnia
- (d) Breathing related sleep disorder
- (e) Primary hypersomnia

136. Hypersomnia is likely to be due to:

- (a) Depressants
- (b) Schizophrenia
- (c) Mania
- (d) Bone fracture
- (e) Anxiety disorder

137. In treatment of insomnia:

- (a) High doses of hypnotics are recommended for most patients
- (b) Dependence rarely occur
- (c) Patient should not know what drug they are on since they are likely to abuse it
- (d) Zolpidem may be used
- (e) There is little or no need for investigations

138. Postpartum psychosis:

- (a) Is an example of psychotic disorder not otherwise specified
- (b) Poses no danger to the patient and others
- (c) There is no relationship between the disorder and Bipolar I Mood disorder
- (d) Hormonal treatment has been found to be effective
- (e) Most women develop the disorder during their third or more delivery

139. A clinical feature of post partum psychosis is:

- (a) Delirium
- (b) Confabulation
- (c) Hypersomnia
- (d) Passivity phenomena
- (e) Delusion

140. A delusion can best be defined as:

- (a) A false belief that meets specific psychological needs
- (b) Perceptual misrepresentation of a sensory image
- (c) Perceptual representation of a sound or object not actually present
- (d) View point able to be changed when convincing evidence to the contrary is presented
- (e) Dissociative reaction

141. Schizophrenia psychotic symptoms such as hallucination, delusion, disorganized speech and grossly disorganized or catatonic behaviour are known as:

- (a) Positive symptoms
- (b) Negative symptoms
- (c) Melting symptoms
- (d) Catastrophic symptoms
- (e) Personalizing symptoms

In order to assume the sick role intentionally produced physical or psychological symptoms are known as:

- (a) Factitious disorder
- (b) Conversion disorder
- (c) Somatization disorder
- (d) Hypochondriasis
- (e) Obsessive disorder

143 Choose the correct answer with regard to DNR/IR:

- (a) Delirium is characterized by a disturbance of consciousness and a change in cognition that develop over a short period of time
- (b) Tremor, myasthenia, in coordination and urinary incontinence are common neurological symptoms
- (c) Abnormalities in mood, perception and behaviour are common psychiatric symptoms
- (d) Symptoms are worse at night
- (e) All of the above are true

144 All of the following are psychotic disorders except:

- (a) Antisocial personality disorder
- (b) Schizophrenia
- (c) Schizoaffective disorder
- (d) Delusional disorder
- (e) Brief psychotic disorder

145 Psychoeducation after an acute psychotic episode should include:

- (a) Information to the patient and primary care providers in the home
- (b) Information about anticipated side effects of medications
- (c) Information on course of illness
- (d) Information on the diagnosis
- (e) All of the above are true

Theme: Hallucinations (For Questions 146 - 150)

- A. Autoscopic
- B. Extracampine
- C. Hygrie
- D. Hypnagogic
- E. Hypnopompic
- F. Kinesthetic
- G. Reflex
- H. Second person auditory
- I. Thalamic

Now decide which delusion that is described in each of the following cases.

146. A patient with a persistent delusion that he is a vampire tells you that he often looks in mirrors and sees no reflection of himself at all. B
147. A schizophrenic gentleman is convinced that he can hear foreign agents in a different city discussing amongst themselves how they are going to assassinate him. D
148. An anxious patient in the ward tells nursing staff how he saw a hooded figure coming to get him as he was about to fall asleep. D
149. A patient becomes highly distressed in response to seeing workmen putting up a shelf on the ward. On questioning she states she can feel them hammering into her head. A
150. An elderly gentleman with psychotic depression tells you that he hears a voice telling him that he is more evil than Hitler and how he should go about killing himself to save the world from his sinfulness. H
