

72. With regard to massive hemothorax the following are true except

- A. 200ml per min in the underwater seal drainage of thoracostomy tube ✓
- B. 1.5l at the connection of chest tube ✓
- C. Always require thoracotomy ✓
- D. 200ml per hour in the underwater seal drainage of thoracostomy tube ✗
- E. May require transfusion ✓

73. The following are forms of primary lymphoedema except

- A. Milroy's disease
- B. Meige's disease
- C. Primary lymphatic malignancy
- D. Lymphoedema Tarda
- E. Sporadic praecox lymphoedema

74. The following features favour late dumping syndrome over early dumping syndrome

- A. At time present with diarrhoea
- B. Presents almost immediately after meal
- C. Relieved by lying down
- D. Aggravated by exercise
- E. Caused by osmotic pressure

75. With regard to gastritis

- A. Type B is autoimmune
- B. Type A is caused by H. Pylori
- C. Type A spares the antrum
- D. Type B leads to pernicious anaemia
- E. Intestinal metaplasia is associated with Type ^B/_A gastritis

76. The following parameters is used in Ranson Criteria within 48 hours after admission

- A. Serum albumin
- B. White blood cell count
- C. Aspartate aminotransferase
- D. Blood urea nitrogen
- E. Blood glucose