

87. Regarding benign breast disease

- A. Cyclical mastalgia is not the commonest reason for referral to the breast clinic
- B. Fibroadenomas are derived from the breast lobule
- C. Lactational breast abscesses are usually due to *Enterobacteriaceae*
- D. Duct ectasia is least common in smokers

E. Atypical lobular hyperplasia is associated with an increased risk of breast cancer

88. A 62 year man presents with gross painless haematuria for the last two months. The following are differential diagnosis except

- A. Kidney stones
- ~~B. Cystitis~~
- C. Urological trauma
- D. Glomerular disease
- E. Hydrocele

89. Regarding varicocele

A. Most are noticed in childhood F

B. The left is affected in 95% of cases

C. In many cases the dilated veins are pampiniform and not cremasteric

D. Does not decompress in supine position

E. Never caused by obstruction of left testicular vein by renal tumour F

90. Regarding hydrocele

A. Uncommon in hot countries F

B. Infantile hydrocele only occur in infancy F

C. About 40% of inguinal hernia are associated with a vaginal hydrocele F

D. Lord's procedure is appropriate in thin walled sac

E. Calcification of the sac is never a complication

* Lord's Procedure
& Jaboulet's procedure

Left cremasteric vein mainly
left testicular vein by renal tumour
Doesn't decompress in supine
position. Affects of testes

Hydro