

47. Indication for laparotomy in the management of sub acute intestinal obstruction due to adhesions are the following except

- A. Rise in pulse
- B. Rise in Respiratory
- C. Two air fluid levels
- D. Rise in WBC over $17 \times 10^9/L$
- E. Increase in ABD Girth

abdomen

48. Which operation in the following surgical disease fall under clean contaminated category

- A. Perforated acute appendicitis (operation of) *dirty*
- B. Perforated duodenal ulcer (operation of)
- C. Repair of inguinal hernia - clean
- D. Elective appendicectomy - clean-contaminated
- E. Colonic perforation - dirty

49. The following are operations for the repair of inguinal hernias in adults except

- A. Bassini repair *No longer used coz of high recurrence rate & abscess*
- B. Shouldice repair
- C. McVay repair
- D. Lichtenstein
- E. Herniotomy

Lyon Method

New repair
not used & intro

Ilip add

50. The most common solid tumor in children in our set up:

- A. Neuroblastoma *
- B. Malignant melanoma
- C. Benign sacrococcygeal teratoma
- D. Malignant sacrococcygeal teratoma *
- E. Embryoma of the kidney

Fernandez
Indirect inguinal
Umbilical

Frequency of
strangulation

Strangulated hernia - Cough impulse is lost
- is irreducible